

**A LA RECHERCHE DE LA TOPOLOGIE
PROJECTIVE. DU CÔTÉ DE CHEZ ARNOLD**

S.YU. OREVKOV

Say that a hypersurface H in $\mathbb{R}\mathbb{P}^n$ is (k, l) -convex or *quasiconvex* if the second fundamental form has the constant signature (k, l) , $k + l = n - 1$. Arnold [1] discovered that the quasiconvexity imposes very strong restrictions on the topology of H . He conjectured the following properties of such H (proven in [1] for $k = 0$):

- 1 (Standardness). The set of all (k, l) -convex hypersurfaces is connected.
- 2 (Covering). $H = S^k \times S^l / (x, y) \sim (-x, -y)$, i.e., H is diffeomorphic to a quadric.
- 3 (Quasistarlikeness). (i). H separates suitable planes $L^+ = \mathbb{R}\mathbb{P}^k$ and $L^- = \mathbb{R}\mathbb{P}^l$ and (ii) any straight line segment $[a, b]$, $a \in L^+$, $b \in L^-$, transversally meets H at a single point.
- 4 (Divisors). For any hyperplane section D of H , the pair (H, D) is diffeomorphic to (\bar{H}, \bar{D}) for a quadric \bar{H} and its suitable hyperplane section \bar{D} .

Hypersurfaces with 3(i) are called *weakly quasistarlike*. The first aim of my talk is to show that none of these conjectures is true for all (k, l) . The second aim is to formulate a similar conjecture which is plausible for any (k, l) .

It is amusing that the simplest counter-example (constructed by E. Cartan in 1938) is closely related to the title of [1]. Namely, let $f : \mathbb{C}\mathbb{P}^2 \rightarrow S^4$ be the double covering constructed in [1] and let $p : S^4 \rightarrow \mathbb{R}\mathbb{P}^4$ be the standard projection. Let H be a tube of constant radius over $p(f(\mathbb{R}\mathbb{P}^2))$. Then H is $(1, 2)$ -convex, but it is not homeomorphic to any quadric. It is not weakly quasistarlike neither.

The Cartan's hypersurface H is *isoparametric*, i.e., its principal curvatures are constant. Any isoparametric hypersurface in $\mathbb{R}\mathbb{P}^n$ is quasiconvex. Isoparametric hypersurfaces in S^n (hence in $\mathbb{R}\mathbb{P}^n$ also) are rather well studied and their classification is almost completed. For $l = k \geq 2$ and for $l = 2k$, $k = 1, 2, 4, 8$, there exist isoparametric (k, l) -convex hypersurfaces in $\mathbb{R}\mathbb{P}^n$ which are not homeomorphic to quadrics (which are not weakly quasistarlike neither). For other values of (k, l) , any (k, l) -convex hypersurface in $\mathbb{R}\mathbb{P}^n$ is a quadric.

It seems plausible that any $(1, 1)$ -convex surface $H \subset \mathbb{R}\mathbb{P}^3$ has properties 1, 2, 3(i), and 4. However, it is easy to construct an example where 3(ii) does not hold. Indeed, let $S^3 \rightarrow \mathbb{R}\mathbb{P}^3 \xrightarrow{h} \mathbb{C}\mathbb{P}^1$ be the Hopf fibration. Set $H = h^{-1}(\gamma)$ where γ is a simple closed path which contains a sufficiently long segment of a spiral.

Conjecture. (Self-Duality). $((\mathbb{R}\mathbb{P}^n)^*, H^*)$ is diffeomorphic to $(\mathbb{R}\mathbb{P}^n, H)$ where H^* is the projectively dual hypersurface of an embedded quasiconvex $H \subset \mathbb{R}\mathbb{P}^n$.

REFERENCES

1. V.I. Arnold, *Ramified covering $CP^2 \rightarrow S^4$, hyperbolicity, and projective topology*, Siberian Math. J. **29:5** (1988), 36-47.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX