

TORUS ACTIONS AND INTEGRABLE SYSTEMS

NGUYEN TIEN ZUNG

ABSTRACT. This is a survey on natural local torus actions which arise in integrable dynamical systems, and their relations with other subjects, including: reduced integrability, local normal forms, affine structures, monodromy, global invariants, integrable surgery, convexity properties of momentum maps, localization formulas, integrable PDEs.

CONTENTS

1. Introduction	2
2. Integrability, torus actions, and reduction	3
2.1. Integrability à la Liouville	3
2.2. Generalized Liouville integrability	4
2.3. Non-Hamiltonian integrability	6
2.4. Reduced integrability of Hamiltonian systems	7
2.5. Non-Hamiltonian reduced integrability	12
3. Torus actions and local normal forms	13
3.1. Toric characterization of Poincaré-Birkhoff normal form	13
3.2. Some simple consequences and generalizations	16
3.3. Convergent normalization for integrable systems	17
3.4. Torus action near a compact singular orbit	20
4. Nondegenerate singularities	22
4.1. Nondegenerate singular points	22
4.2. Nondegenerate singular orbits	23
4.3. Nondegenerate singular fibers	25
4.4. Focus-focus singularities	26
5. Global aspects of local torus actions	27
5.1. Sheaf of local \mathbb{T}^1 -actions	27
5.2. Affine base space, integrable surgery, and convexity	28
5.3. Localization formulas	30
6. Infinite-dimensional torus actions	30
References	32

Date: second version, July 2004.

1991 Mathematics Subject Classification. 37J35,53D20,37G05,70K45,34C14.

Key words and phrases. integrable system, local torus actions, Poincaré-Birkhoff normal form, reduced integrability, affine structure, monodromy, convexity, proper groupoid, localization formula.

1. INTRODUCTION

To say that everything is a torus would be a great exaggeration, but to say that *everything contains a torus* would not be too far from the truth. According to ancient oriental philosophy, everything can be described by (a combination of) five elemental aspects, or phases: regular, transitive, expansive, chaotic, and contractive, and if we look at these five phases as a whole then they also form cycles.

This survey paper is concerned with regular aspects of things. Mathematically, they correspond to regular dynamics, or integrable dynamical systems. For me, *an integrable system is a local torus action*. The main dynamical property of a regular dynamical system is its quasi-periodic behavior. Mathematically, it means that locally there is a torus action which preserves the system. These torus actions exist near compact regular orbits (Liouville's theorem). To a great extent, they exist near singularities of integrable systems as well, and the main topics of this paper are how to find them and what are the implications of their existence.

I spent the last fifteen years looking for tori and this paper is mostly a report on what I found – probably very little for a fifteen year work. My lame excuse is that I often had an empty stomach (empty pocket) to the point of wanting to forget completely about the tori on more than one occasions. This “toric business” began in 1989 when I was a 3rd year undergraduate student under the direction of A.T. Fomenko: for my year-end memoir I studied integrable perturbations of integrable Hamiltonian systems of 2 degrees of freedom, and found out that there is a local Hamiltonian \mathbb{T}^1 -action which preserves an integrable 2D system near each corank 1 nondegenerate singular hyperbolic level set [91]. This little discovery is the starting point for more general existence results obtained later.

The topics discussed in this paper can be seen from the table of contents. They include: reduced integrability, Poincaré–Birkhoff normalization, automorphism groups, partial action-angle variables, classification of singularities, monodromy, characteristic classes, integrable surgery, convexity of momentum maps, localization formulas, and so on. We consider only classical dynamical systems in this paper. It turns out that many local and semi-local results about the behavior of classical integrable systems have their counterparts in quantum integrable systems, or at least are useful for the study of quantum systems, see Vu Ngoc San [76] and references therein.

The present paper deals mainly with finite-dimensional dynamical systems, i.e. ordinary differential equations, though in the last section we will briefly discuss the infinite dimensional case, i.e. integrable PDEs, where there is a huge amount of literature but at the same time many basic questions on topological aspects remain open.

Acknowledgements. I would like to thank my Russian colleagues Alexei Bolsinov, Anatoly Fomenko and Andrey Oshemkov for the invitation to write this paper.

Many people helped me find the tori. I'm indebted to them all, and especially to Michèle Audin, Alexei Bolsinov, Yves Colin de Verdière, Richard Cushman, Jean-Paul Dufour, Hans Duistermaat, Anatoly Fomenko, Lubomir Gavrilov, Thomas Kappeler, Pierre Molino, Tudor Ratiu, Jean-Claude Sikorav, Vu Ngoc San, and Alan Weinstein.

A part of this paper was written during the author's visit to Bernoulli Center, EPFL, Lausanne, and he would like to thank Tudor Ratiu and the Bernoulli Center for the hospitality.

2. INTEGRABILITY, TORUS ACTIONS, AND REDUCTION

2.1. Integrability à la Liouville.

Probably the most well-known notion of integrability in dynamical systems is the notion of integrability à la Liouville for Hamiltonian systems on symplectic manifolds. Denote by (M^{2n}, ω) a symplectic manifold of dimension $2n$ with symplectic form ω , and H a function on M^{2n} . Denote by X_H the Hamiltonian vector field of H on M^{2n} :

$$(2.1) \quad i_{X_H}\omega = -dH .$$

Definition 2.1. A function H (or the corresponding Hamiltonian vector field X_H) on a $2n$ -dimensional symplectic manifold (M^{2n}, ω) is called **integrable à la Liouville**, or **Liouville-integrable**, if it admits n functionally independent first integrals in involution. In other words, there are n functions $F_1 = H, F_2, \dots, F_n$ on M^{2n} such that $dF_1 \wedge \dots \wedge dF_n \neq 0$ almost everywhere and $\{F_i, F_j\} = 0 \forall i, j$.

In the above definition, $\{F_i, F_j\} := X_{F_i}(F_j)$ denotes the Poisson bracket of F_i and F_j with respect to the symplectic form ω . The map

$$(2.2) \quad \mathbf{F} = (F_1, \dots, F_n) : (M^{2n}, \omega) \rightarrow \mathbb{K}^n$$

is called the **momentum map** ($\mathbb{K} = \mathbb{R}$ or \mathbb{C}). The above definition works in many categories: smooth, real analytic, holomorphic, formal, etc.

The condition $X_H(F_i) = \{H, F_i\} = 0$ implies that the Hamiltonian vector field X_H is tangent to the level sets of \mathbf{F} . Let $N = \mathbf{F}^{-1}(c)$ be a regular connected (component of a) level set of \mathbf{F} . Then it is a Lagrangian submanifold of M^{2n} : the dimension of N is half the dimension of M^{2n} , and the restriction of ω to N is zero. So we can talk about a (singular) Lagrangian foliation/fibration given by the momentum map.

A classical result attributed to Liouville [58] says that, in the smooth case, if a connected level set N is compact and does not intersect with the boundary of M^{2n} , then it is diffeomorphic to a standard torus \mathbb{T}^n , and the Hamiltonian system X_H is quasi-periodic on N : in other words, there is a periodic coordinate system (q_1, \dots, q_n) on N with respect to which the restriction of X_H to N has constant coefficients: $X_H = \sum \gamma_i \partial / \partial q_i$, γ_i being constants. For this reason, N is called a **Liouville torus**.

The description of a Liouville-integrable Hamiltonian system near a Liouville torus is given by the following theorem about the existence of action-angle variables. This theorem is often called Arnold-Liouville theorem, but it was essentially obtained by Henri Mineur in 1935 [63, 64]:

Theorem 2.2 (Liouville–Mineur–Arnold). *Let N be a Liouville torus of a Liouville-integrable Hamiltonian system with a given momentum map $\mathbf{F} : (M^{2n}, \omega) \rightarrow \mathbb{R}^n$.*

Then there is a neighborhood $\mathcal{U}(N)$ of N and a smooth symplectomorphism

$$(2.3) \quad \Psi : (\mathcal{U}(N), \omega) \rightarrow (D^n \times \mathbb{T}^n, \sum_1^n d\nu_i \wedge d\mu_i)$$

(ν_i - coordinates of D^n , $\mu_i \pmod{1}$ - periodic coordinates of \mathbb{T}^n) such that \mathbf{F} depends only on $I_i = \phi^* \nu_i$, i.e. \mathbf{F} does not depend on $\phi_i = \phi^* \mu_i$.

The variables (I_i, ϕ_i) in the above theorem are called **action-angle variables**. The map

$$(2.4) \quad (I_1, \dots, I_n) : (\mathcal{U}(N), \omega) \rightarrow \mathbb{R}^n$$

is the momentum map of a Hamiltonian torus \mathbb{T}^n -action on $(\mathcal{U}(N), \omega)$ which preserves \mathbf{F} . The existence of this Hamiltonian torus action is essentially equivalent to Liouville-Mineur-Arnold theorem: once the action variables are found, angle variables can also be found easily by fixing a Lagrangian section to the foliation by Liouville tori. The quasi-periodicity of the system on N also follows immediately from the existence of this torus action.

The existence of action-angle variables is very important, both for the theory of near-integrable systems (K.A.M. theory), and for the quantization of integrable systems (Bohr–Sommerfeld rule). Actually, Mineur was an astrophysicist, and Bohr–Sommerfeld quantization was his motivation for finding action-angle variables.

Mineur [64] also wrote down the following simple formula, which we will call **Mineur–Arnold formula**, for action functions:

$$(2.5) \quad I_i(z) = \int_{\Gamma_i(z)} \beta$$

where z is a point in $\mathcal{U}(N)$, β is a primitive of the symplectic form ω , i.e. $d\beta = \omega$, and $\Gamma_i(z)$ is an 1-cycle on the Liouville torus which contains z (and which depends on z continuously).

In the case of *algebraically integrable systems* (see e.g. [1]), where invariant tori can be identified with (the real part of) Jacobian or Prym varieties of complex curves (*spectral curves* of the system), the integral in Mineur–Arnold formula corresponds to Abelian integrals on complex curves, as observed by Novikov and Veselov [81].

It often happens that the above Mineur–Arnold formula (2.5) is still valid when the cycle Γ_i lies on a singular level set, and it leads to an action function near a singularity (see e.g. [36, 43] and Section 3).

2.2. Generalized Liouville integrability.

In practice, one often deals with Hamiltonian systems which admit a non-Abelian group of symmetries, or Hamiltonian systems on Poisson (instead of symplectic) manifolds. A typical example is the Euler equation on the dual of a Lie algebra. For such systems, Liouville integrability needs to be replaced by a more general and convenient notion of integrability, which nevertheless retains the main feature of Liouville integrability, namely the existence of local torus actions.

Let (M, Π) be a **Poisson manifold**, with Π being the Poisson structure. It means that Π is a 2-vector field on M such that the following binary operation on

the space of functions on M , called the **Poisson bracket**,

$$(2.6) \quad \{H, F\} = \langle dH \wedge dF, \Pi \rangle$$

is a Lie bracket, i.e. it satisfies the Jacobi identity. A symplectic manifold is also a Poisson manifold. Conversely, a Poisson manifold can be seen as a singular foliation by symplectic manifolds, see e.g. [85].

Let H be a function on a Poisson manifold (M, Π) , and X_H the corresponding Hamiltonian vector field: $X_H = dH \lrcorner \Pi$. Let \mathcal{F} be a set of first integrals of X_H , i.e. each $F \in \mathcal{F}$ is a function on M which is preserved by X_H (equivalently, $\{F, H\} = 0$). Denote by $\text{ddim } \mathcal{F}$ the **functional dimension** of \mathcal{F} , i.e. the maximal number of functions in \mathcal{F} which are functionally independent almost everywhere.

We will associate to \mathcal{F} the space $\mathcal{X}_{\mathcal{F}}$ of Hamiltonian vector fields X_E such that $X_E(F) = 0$ for all $F \in \mathcal{F}$ and E is functionally dependent of \mathcal{F} (i.e. the functional dimension of the union of \mathcal{F} with the function E is the same as the functional dimension of \mathcal{F}). Clearly, the vector fields in $\mathcal{X}_{\mathcal{F}}$ commute pairwise and commute with X_H . Denote by $\text{ddim } \mathcal{X}_{\mathcal{F}}$ the functional dimension of $\mathcal{X}_{\mathcal{F}}$, i.e. the maximal number of vector fields in $\mathcal{X}_{\mathcal{F}}$ which are linearly independent at almost every point. Note that we always have $\text{ddim } \mathcal{F} + \text{ddim } \mathcal{X}_{\mathcal{F}} \leq m$, because the vector fields in $\mathcal{X}_{\mathcal{F}}$ are tangent to the common level sets of the functions in \mathcal{F} .

The following definition is essentially due to Nekhoroshev [68] and Mischenko and Fomenko [67] :

Definition 2.3. A Hamiltonian vector field X_H on an m -dimensional Poisson manifold (M, Π) is called **integrable in generalized Liouville sense** if there is a set of first integrals \mathcal{F} such that $\text{ddim } \mathcal{F} + \text{ddim } \mathcal{X}_{\mathcal{F}} = m$.

The above notion of integrability is also called **noncommutative integrability**, due to the fact that the functions in \mathcal{F} do not Poisson-commute in general, and in many cases one may choose \mathcal{F} to be a finite-dimensional non-commutative Lie algebra of functions under the Poisson bracket. When the functions in \mathcal{F} Poisson-commute and the Poisson structure is nondegenerate, we get back to the usual integrability à la Liouville.

Denote $q = \text{ddim } \mathcal{F}, p = \text{ddim } \mathcal{X}_{\mathcal{F}}$. Then we can find p Hamiltonian vector fields $X_1 = X_{E_1}, \dots, X_p = X_{E_p} \in \mathcal{X}_{\mathcal{F}}$ and q functions $F_1, \dots, F_q \in \mathcal{F}$ such that we have:

$$(2.7) \quad \begin{aligned} X_H(F_i) = 0, [X_H, X_i] = 0, [X_i, X_j] = 0, X_i(F_j) = 0 \quad \forall i, j, \\ X_1 \wedge \dots \wedge X_p \neq 0 \text{ and } dF_1 \wedge \dots \wedge dF_q \neq 0 \text{ almost everywhere.} \end{aligned}$$

The existence of such a p -tuple $\mathbf{X} = (X_1, \dots, X_p)$ of commuting Hamiltonian vector fields and q -tuple $\mathbf{F} = (F_1, \dots, F_q)$ of common first integrals with $p + q = m$ is equivalent to the integrability in the generalized Liouville sense. When $p + q = m$, we will say that H is integrable with the aid of (\mathbf{X}, \mathbf{F}) , and by abuse of language, we will also say that (\mathbf{X}, \mathbf{F}) is an **integrable Hamiltonian system** in generalized Liouville sense. The map

$$(2.8) \quad \mathbf{F} = (F_1, \dots, F_q) : (M, \Pi) \rightarrow \mathbb{K}^q$$

(where $\mathbb{K} = \mathbb{R}$ or \mathbb{C}) is called the **generalized momentum map**. The (regular) level sets of this map are called **invariant manifolds**: they are invariant with respect to X_H, \mathbf{X} and \mathbf{F} . They are of dimension p , lie on the symplectic leaves of M , and are isotropic. When $p < \frac{1}{2} \text{rank } \Pi$, i.e. when the invariant manifolds

are isotropic but not Lagrangian, one also speaks of **degenerate integrability**, or **superintegrability**, see e.g. [32, 68, 71].

Definition 2.4. With the above notations, a Hamiltonian system X_H , on a real Poisson manifold (M, Π) , integrable with the aid of (\mathbf{X}, \mathbf{F}) , is called **proper** if the generalized momentum map $\mathbf{F} : M \rightarrow \mathbb{R}^q$ is a proper map from M to its image, and the image of the singular set $\{x \in M, X_1 \wedge X_2 \wedge \dots \wedge X_p(x) = 0\}$ of the commuting Hamiltonian vector fields under the momentum map $\mathbf{F} : M \rightarrow \mathbb{R}^q$ is nowhere dense in \mathbb{R}^q .

Under the properness condition, one gets a natural generalization of the classical Liouville-Mineur-Arnold theorem [68, 67]: outside the singular region, the Poisson manifold M is foliated by invariant isotropic p -dimensional tori on which the flow of X_H is quasi-periodic, and there exist local action-angle coordinates. The action variables can still be defined by Mineur-Arnold formula (2.5). There will be p action and p angle variables (so one will have to add $(q - p)$ variables to get a full system of variables). In particular, near every isotropic invariant torus there is a free Hamiltonian \mathbb{T}^p -action which preserves the system.

Example 2.5. A Hamiltonian \mathbb{T}^p -action on a Poisson manifold can be seen as a proper integrable system – the space of first integrals is the space of \mathbb{T}^p -invariant functions, and in this case we have a *global* \mathbb{T}^p -action which preserves the system. More generally, one can associate to each Hamiltonian action of a compact Lie group G on a Poisson manifold a proper integrable system: H is the composition of the momentum map $\mu : M \rightarrow \mathfrak{g}^*$ with a generic Ad^* -invariant function $h : \mathfrak{g}^* \rightarrow \mathbb{R}$, see Subsection 2.4.

Remark 2.6. There is a natural question: is an integrable Hamiltonian system in generalized Liouville sense on a symplectic manifold also integrable à la Liouville? One often expects the answer to be Yes. See e.g. Fomenko [34] for a long discussion on this question, and the related question about the existence of Liouville-integrable systems on given symplectic manifolds.

Remark 2.7. Another natural question is the following. Let \mathcal{F}_H denote the space of all first integrals of H . Suppose that H is integrable in generalized Liouville sense. Is it true that H is integrable with the aid of $(\mathcal{F}_H, \mathcal{X}_{\mathcal{F}_H})$, i.e. $\dim \mathcal{F}_H + \dim \mathcal{X}_{\mathcal{F}_H} = 0$? We expect the answer to be yes for “reasonable” systems. It is easy to see that the answer is Yes in the proper integrable case, under the additional assumption that the orbits of X_H are dense (i.e. its frequencies are incommensurable) on almost every invariant torus (i.e. common level of a given set of first integrals \mathcal{F}). In this case $\mathcal{X}_{\mathcal{F}_H}$ consists of the Hamiltonian vector fields whose flow is quasi-periodic on each invariant torus. Another case where the answer is also Yes arises in the study of local normal forms of analytic integrable vector fields, see Section 3.

2.3. Non-Hamiltonian integrability.

There are many physical non-Hamiltonian (e.g. non-holonomic) systems, which may naturally be called integrable in a non-Hamiltonian sense, because their behavior is very similar to that of integrable Hamiltonian systems, see e.g. [7, 22]. A simple example is the Chinese top. (It is a spinning top whose lower part looks like a hemisphere and whose upper part is heavy. When you spin it, it will turn upside

down after a while). The notion of non-Hamiltonian integrability was probably first introduced by Bogoyavlenskij (see [10] and references therein), who calls it **broad integrability**, though other authors also arrived at it independently, from different points of view, see e.g. [7, 10, 22, 78, 96].

Definition 2.8. A vector field X on a manifold M is called **integrable in non-Hamiltonian sense** with the aid of $(\mathcal{F}, \mathcal{X})$, where \mathcal{F} is a set of functions on M and \mathcal{X} is a set of vector fields on M , if the following conditions are satisfied :

- a) $X(F) = 0$ and $Y(F) = 0 \ \forall \ F \in \mathcal{F}, Y \in \mathcal{X}$,
- b) $[Y, X] = [Y, Z] = 0 \ \forall \ Y, Z \in \mathcal{X}$,
- d) $\dim M = \text{ddim } \mathcal{F} + \text{ddim } \mathcal{X}$.

In the real case, if, moreover, there is a p -tuple $\mathbf{X} = (X_1, \dots, X_p)$ of vector fields in \mathcal{X} and a q -tuple $\mathbf{F} = (F_1, \dots, F_q)$ of functionally independent functions in \mathcal{F} , where $p = \text{ddim } \mathcal{X}$ and $q = \text{ddim } \mathcal{F}$, such that the map $\mathbf{F} : M \rightarrow \mathbb{R}^q$ is a proper map from M to its image, and for almost every level set of this map the vector fields X_1, \dots, X_p are linearly independent everywhere on this level set, then we say that X is **proper integrable** with the aid of (\mathbf{X}, \mathbf{F}) , and by abuse of language we will also say that (\mathbf{X}, \mathbf{F}) is a proper integrable non-Hamiltonian system of bi-degree (p, q) of freedom.

So non-Hamiltonian integrability is almost the same as Hamiltonian integrability, except for the fact that the vector fields X, X_1, \dots, X_p are not required to be Hamiltonian. It is not surprising that Liouville's theorem holds for proper non-Hamiltonian integrable systems as well: each regular invariant manifold (connected level set of \mathbf{F}) is a p -dimensional torus on which the system is quasi-periodic, and in a neighborhood of it there is a free \mathbb{T}^p -torus action which preserves the system.

If a Hamiltonian system is (proper) integrable in the generalized Liouville sense, then of course it is also (proper) integrable in the non-Hamiltonian sense, though the inverse is not true: it may happen that the invariant tori are not isotropic, see e.g. [10].

Remark 2.9. Remark 2.7 also applies to non-Hamiltonian systems: For an integrable vector field X on a manifold M , denote by \mathcal{F}_X the set of all first integrals of X , and by \mathcal{X}_X the set of all vector fields which commute with X and preserve every function in \mathcal{F} . Then a natural question is, do we have the equality $\text{ddim } \mathcal{F}_X + \text{ddim } \mathcal{X}_X = \dim M$? The answer is similar to the Hamiltonian case. In particular, if the system is proper and the vector field X is nonresonant (i.e. has a dense orbit) on almost every invariant torus, then the answer is yes.

Remark 2.10. If (\mathbf{X}, \mathbf{F}) is a non-Hamiltonian integrable system of bidegree (p, q) on a manifold M , then it can be lifted to a Liouville-integrable system on the cotangent bundle T^*M . Denote by $\pi : T^*M \rightarrow M$ the projection, then the corresponding momentum map is $(H_1, \dots, H_{p+q}) : T^*M \rightarrow \mathbb{R}^{p+q}$, where $H_i = \pi^* F_i$ ($i = 1, \dots, q$), and $H_{i+q}(\alpha) = \langle \alpha, X_i(\pi(\alpha)) \rangle \ \forall \ \alpha \in T^*M$ ($i = 1, \dots, p$).

2.4. Reduced integrability of Hamiltonian systems.

In the literature, when people speak about integrability of a dynamical system, they often actually mean its **reduced integrability**, i.e. integrability of the reduced (with respect to a natural symmetry group action) system. For example, consider an integrable spinning top (e.g. the Kovalevskaya top). Its configuration

space is $SO(3)$, so it is naturally a Hamiltonian system with 3 degrees of freedom. But it is often considered as a 2-degree-of-freedom integrable system with a parameter, see e.g. [11].

Curiously, to my knowledge, the natural question about the effect of reduction on integrability has never been formally addressed in monographs on dynamical systems. Recently we studied this question [97], and showed that, for a Hamiltonian system invariant under a proper action of a Lie group, integrability is essentially equivalent to reduced integrability.

It turns out that the most natural notion of integrability to use here is not the Liouville integrability, but rather the integrability in generalized Liouville sense. Also, since the category of manifolds is not invariant under the operation of taking quotient with respect to a proper group action, we have to replace manifolds by generalized manifolds: in this paper, a **generalized manifold** is a differentiable space which is locally isomorphic to the quotient of a manifold by a compact group action. Due to well-known results about functions invariant under compact group actions, see e.g. [70], one can talk about smooth functions, vector fields, differential forms, etc. on generalized manifolds, and the previous integrability definitions work for them as well.

Let (M, Π) be a Poisson generalized manifold, G a Lie group which acts properly on M , H a function on M which is invariant under the action of G . Then the quotient space M/G is again a Poisson generalized manifold, see e.g. [23]. We will denote the projection of Π, H, X_H on M/G by $\Pi/G, H/G, X_H/G$ respectively. Of course, X_H/G is the Hamiltonian vector field of H/G .

We will assume that the action of G on (M, Π) is Hamiltonian, with an equivariant moment map $\mu : M \rightarrow \mathfrak{g}^*$, where \mathfrak{g} denotes the Lie algebra of G , and that the following additional condition is satisfied: Recall that the image $\mu(M)$ of M under the moment map $\mu : M \rightarrow \mathfrak{g}^*$ is saturated by symplectic leaves (i.e. coadjoint orbits) of \mathfrak{g}^* . Denote by s the minimal codimension in \mathfrak{g}^* of a coadjoint orbit which lies in $\pi(M)$. Then the additional condition is that there exist s functions f_1, \dots, f_s on \mathfrak{g}^* , which are invariant on the coadjoint orbits which lie in $\mu(M)$, and such that for almost every point $x \in M$ we have $df_1 \wedge \dots \wedge df_s(\mu(x)) \neq 0$. For example, when G is compact and M is connected, then this condition is satisfied automatically.

With the above notations and assumptions, we have :

Theorem 2.11 ([97]). *If the system $(M/G, X_H/G)$ is integrable in generalized Liouville sense, then the system (M, X_H) also is. Moreover, if G is compact and $(M/G, X_H/G)$ is proper, then (M, X_H) also is.*

Since the preprint [97] will not be published as a separate paper, let us include here a full proof of Theorem 2.11.

Proof. Denote by \mathcal{F}' a set of first integrals of X_H/G on M/G which provides the integrability of X_H/G , and by $\mathcal{X}' = \mathcal{X}_{\mathcal{F}'}$ the corresponding space of commuting Hamiltonian vector fields on M/G . We have $\dim M/G = p' + q'$ where $p' = \text{ddim } \mathcal{X}'$ and $q' = \text{ddim } \mathcal{F}'$.

Recall that, by our assumptions, there exist s functions f_1, \dots, f_s on \mathfrak{g}^* , which are functionally independent almost everywhere in $\mu(M)$, and which are invariant on the coadjoint orbits which lie in $\mu(M)$. Here s is the minimal codimension in

\mathfrak{g}^* of the coadjoint orbits which lie in $\mu(M)$. We can complete (f_1, \dots, f_s) to a set of d functions $f_1, \dots, f_s, f_{s+1}, \dots, f_d$ on \mathfrak{g}^* , where $d = \dim G = \dim \mathfrak{g}$ denotes the dimension of \mathfrak{g} , which are functionally independent almost everywhere in $\mu(M)$.

Denote by $\overline{\mathcal{F}}$ the pull-back of \mathcal{F}' under the projection $\mathfrak{p} : M \rightarrow M/G$, and by F_1, \dots, F_d the pull-back of f_1, \dots, f_d under the moment map $\mu : M \rightarrow \mathfrak{g}^*$. Note that, since H is G -invariant, the functions F_i are first integrals of X_H . And of course, $\overline{\mathcal{F}}$ is also a set of first integrals of X_H . Denote by \mathcal{F} the union of $\overline{\mathcal{F}}$ with (F_{s+1}, \dots, F_d) . (It is not necessary to include F_1, \dots, F_s in this union, because these functions are G -invariant and project to Casimir functions on M/G , which implies that they are functionally dependent of $\overline{\mathcal{F}}$). We will show that X_H is integrable with the aid of \mathcal{F} .

Notice that, by assumptions, the coadjoint orbits of \mathfrak{g}^* which lie in $\mu(M)$ are of generic dimension $d-s$, and the functions f_{s+1}, \dots, f_d may be viewed as a coordinate system on a symplectic leaf of $\mu(M)$ at a generic point. In particular, we have

$$\langle df_{s+1} \wedge \dots \wedge df_d, X_{f_{s+1}} \wedge \dots \wedge X_{f_d} \rangle \neq 0,$$

which implies, by equivariance :

$$\langle dF_{s+1} \wedge \dots \wedge dF_d, X_{F_{s+1}} \wedge \dots \wedge X_{F_d} \rangle \neq 0.$$

Since the vector fields $X_{F_{s+1}}, \dots, X_{F_d}$ are tangent to the orbits of G on M , and the functions in $\overline{\mathcal{F}}$ are invariant on the orbits of G , it implies that the set (F_{s+1}, \dots, F_d) is “totally” functionally independent of $\overline{\mathcal{F}}$. In particular, we have :

$$(2.9) \quad \text{ddim } \mathcal{F} = \text{ddim } \mathcal{F}' + \text{ddim } (F_{s+1}, \dots, F_d) = q' + d - s,$$

where $q' = \text{ddim } \mathcal{F}'$. On the other hand, we have

$$\dim M = \dim M/G + (d - k) = p' + q' + d - k,$$

where $p' = \text{ddim } \mathcal{X}_{\mathcal{F}'}$, and k is the dimension of a minimal isotropic group of the action of G on M . Thus, in order to show the integrability condition

$$\dim M = \text{ddim } \mathcal{F} + \text{ddim } \mathcal{X}_{\mathcal{F}},$$

it remains to show that

$$(2.10) \quad \text{ddim } \mathcal{X}_{\mathcal{F}} = \text{ddim } \mathcal{X}_{\mathcal{F}'} + (s - k).$$

Consider the vector fields $Y_1 = X_{F_1}, \dots, Y_d = X_{F_d}$ on M . They span the tangent space to the orbit of G on M at a generic point. The dimension of such a generic tangent space is $d - k$. It implies that, among the first s vector fields, there are at least $s - k$ vector fields which are linearly independent at a generic points : we may assume that $Y_1 \wedge \dots \wedge Y_{s-k} \neq 0$.

Let $X_{h_1}, \dots, X_{h_{p'}}$ be p' linearly independent (at a generic point) vector fields which belong to $\mathcal{X}_{\mathcal{F}'}$, where $p' = \text{ddim } \mathcal{X}_{\mathcal{F}'}$. Then we have

$$X_{\mathfrak{p}^*(h_1)}, \dots, X_{\mathfrak{p}^*(h_{p'})}, Y_1, \dots, Y_{s-k} \in \mathcal{X}_{\mathcal{F}},$$

and these $p' + s - k$ vector fields are linearly independent at a generic point. (Recall that, at each point $x \in M$, the vectors $Y_1(x), \dots, Y_{s-k}(x)$ are tangent to the orbit of G which contains x , while the linear space spanned by $X_{\mathfrak{p}^*(h_1)}, \dots, X_{\mathfrak{p}^*(h_{p'})}$ contains no tangent direction to this orbit).

Thus we have $\text{ddim } \mathcal{X}_{\mathcal{F}} \geq p' + s - k$, which means that $\text{ddim } \mathcal{X}_{\mathcal{F}} = p' + s - k$ (because, as discussed earlier, we always have $\text{ddim } \mathcal{F} + \text{ddim } \mathcal{X}_{\mathcal{F}} \leq \dim M$). We have proved that if $(M/G, X_H/G)$ is integrable in generalized Liouville sense then (M, X_H) also is.

Now suppose that G is compact and $(M/G, X_H/G)$ is proper: there are q' functionally independent functions $g_1, \dots, g_{q'} \in \mathcal{F}'$ such that $(g_1, \dots, g_{q'}) : M/G \rightarrow \mathbb{R}^{q'}$ is a proper map from M/G to its image, and p' Hamiltonian vector fields $X_{h_1}, \dots, X_{h_{p'}}$ in \mathcal{X}' such that on a generic common level set of $(g_1, \dots, g_{q'})$ we have that $X_{h_1} \wedge \dots \wedge X_{h_{p'}}$ does not vanish anywhere. Then it is straightforward that

$$\mathfrak{p}^*(g_1), \dots, \mathfrak{p}^*(g_{q'}), F_{s+1}, \dots, F_d \in \mathcal{F}$$

and the map

$$(\mathfrak{p}^*(g_1), \dots, \mathfrak{p}^*(g_{q'}), F_{s+1}, \dots, F_d) : M \rightarrow \mathbb{R}^{q'+d-s}$$

is a proper map from M to its image. More importantly, on a generic level set of this map we have that the $(q' + s - k)$ -vector $X_{\mathfrak{p}^*(h_1)} \wedge \dots \wedge X_{\mathfrak{p}^*(h_{p'})} \wedge Y_1 \wedge \dots \wedge Y_{s-k}$ does not vanish anywhere. To prove this last fact, notice that $X_{\mathfrak{p}^*(h_1)} \wedge \dots \wedge X_{\mathfrak{p}^*(h_{p'})} \wedge Y_1 \wedge \dots \wedge Y_{s-k}(x) \neq 0$ for a point $x \in M$ if and only if $X_{\mathfrak{p}^*(h_1)} \wedge \dots \wedge X_{\mathfrak{p}^*(h_{p'})}(x) \neq 0$ and $Y_1 \wedge \dots \wedge Y_{s-k}(x) \neq 0$ (one of these two multi-vectors is transversal to the G -orbit of x while the other one ‘‘lies on it’’), and that these inequalities are $G \times \mathbb{R}^{p'}$ -invariant properties, where the action of $\mathbb{R}^{p'}$ is generated by $X_{\mathfrak{p}^*(h_1)}, \dots, X_{\mathfrak{p}^*(h_{p'})}$. \square

Remark 2.12. Similar results to Theorem 2.11 have been obtained independently by Bolsinov and Jovanovic [12, 46], who used them to construct new examples of integrable geodesic flows, e.g. on biquotients of compact Lie groups.

Example 2.13. The simplest example which shows an evident relationship between reduction and integrability is the classical Euler top : it can be written as a Hamiltonian system on $T^*SO(3)$, invariant under a natural Hamiltonian action of $SO(3)$, is integrable with the aid of a set of four first integrals, and has 2-dimensional isotropic invariant tori. The geodesic flow of a bi-invariant metric on a compact Lie group is also properly integrable : in fact, the corresponding reduced system is trivial (identically zero). More generally, let $H = h \circ \mu$ be a collective Hamiltonian in the sense of Guillemin–Sternberg (see e.g. [40]), where $\mu : M \rightarrow \mathfrak{g}^*$ is the momentum map of a Hamiltonian compact group action, and h is a function on \mathfrak{g}^* . If h is a Casimir function on \mathfrak{g}^* , then H is integrable because its reduction will be a trivial Hamiltonian system.

Remark 2.14. Recall from Equation (2.10) that $\text{ddim } \mathcal{X}_{\mathcal{F}} - \text{ddim } \mathcal{X}_{\mathcal{F}'} = s - k$, where k is the dimension of a generic isotropic group of the G -action on M , and s is the (minimal) corank in \mathfrak{g}^* of a coadjoint orbit which lies in $\mu(M)$. On the other hand, the difference between the rank of the Poisson structure on M and the reduced Poisson structure on M/G can be calculated as follows :

$$(2.11) \quad \text{rank } \Pi - \text{rank } \Pi/G = (d - k) + (s - k)$$

Here $(d - k)$ is the difference between $\dim M$ and $\dim M/G$, and $(s - k)$ is the difference between the corank of Π/G in M/G and the corank of Π in M . It follows that

$$(2.12) \quad \text{rank } \Pi - 2\text{ddim } \mathcal{X}_{\mathcal{F}} = \text{rank } \Pi/G - 2\text{ddim } \mathcal{X}_{\mathcal{F}'} + (d - s)$$

In particular, if $d - s > 0$ (typical situation when G is non-Abelian), then we always have $\text{rank } \Pi - 2\text{ddim } \mathcal{X}_{\mathcal{F}} > 0$ (because we always have $\text{rank } \Pi/G - 2\text{ddim } \mathcal{X}_{\mathcal{F}'} \geq 0$ due to integrability), i.e. the original system is always super-integrable with the aid of \mathcal{F} . When G is Abelian (implying $d = s$), and the reduced system is Liouville-integrable with the aid of \mathcal{F}' (i.e. $\text{rank } \Pi/G = 2\text{ddim } \mathcal{X}_{\mathcal{F}'}$), then the original system is also Liouville-integrable with the aid of \mathcal{F} .

Remark 2.15. Following Mischenko-Fomenko [67], we will say that a hamiltonian system (M, Π, X_H) is **non-commutatively integrable in the restricted sense** with the aid of \mathcal{F} , if \mathcal{F} is a finite-dimensional Lie algebra under the Poisson bracket and (M, Π, X_H) is integrable with the aid of \mathcal{F} . In other words, we have an equivariant moment maps $(M, \Pi) \rightarrow \mathfrak{f}^*$, where \mathfrak{f} is some finite-dimensional Lie algebra, and if we denote by f_1, \dots, f_n the components of this moment map, then they are first integrals of X_H , and X_H is integrable with the aid of this set of first integrals. Theorem 2.11 remains true, and its proof remains the same if not easier, if we replace Hamiltonian integrability by non-commutative integrability in the restricted sense. Indeed, if $M \rightarrow \mathfrak{g}^*$ is the equivariant moment map of the symmetry group G , and if $M/G \rightarrow \mathfrak{h}^*$ is an equivariant moment map which provides non-commutative integrability in the restricted sense on M/G , then the map $M \rightarrow \mathfrak{h}^*$ (which is the composition $M \rightarrow M/G \rightarrow \mathfrak{h}^*$) is an equivariant moment map which commutes with $M \rightarrow \mathfrak{g}^*$, and the direct sum of this two maps, $M \rightarrow \mathfrak{f}^*$ where $\mathfrak{f} = \mathfrak{g} \oplus \mathfrak{h}$, will provide non-commutative integrability in the restricted sense on M .

Theorem 2.11 has the following inverse (see Remark 2.7):

Theorem 2.16. *If G is compact, and if the Hamiltonian system (M, X_H) is integrable with the aid of \mathcal{F}_H (the set of all first integrals of H) in the sense that $\text{ddim } \mathcal{F}_H + \text{ddim } \mathcal{X}_{\mathcal{F}_H} = \dim M$, then the reduced Hamiltonian system $(M/G, X_H)$ is also integrable. Moreover, if (M, X_H) is proper then $(M/G, X_H)$ also is.*

Proof. By assumptions, we have $\dim M = p + q$, where $q = \text{ddim } \mathcal{F}_H$ and $p = \text{ddim } \mathcal{X}_{\mathcal{F}_H}$, and we can find p first integrals H_1, \dots, H_p of H such that X_{H_1}, \dots, X_{H_p} are linearly independent (at a generic point) and belong to $\mathcal{X}_{\mathcal{F}_H}$. In particular, we have $X_{H_i}(F) = 0$ for any $F \in \mathcal{F}$ and $1 \leq i \leq p$.

An important observation is that the functions H_1, \dots, H_p are G -invariant. In deed, if we denote by F_1, \dots, F_d the components of the equivariant moment map $\pi : M \rightarrow \mathfrak{g}^*$ (via an identification of \mathfrak{g}^* with \mathbb{R}^d), then since H is G -invariant we have $\{H, F_j\} = 0$, i.e. $F_j \in \mathcal{F}_H$, which implies that $\{F_j, H_i\} = 0 \ \forall 1 \leq i \leq d, \ 1 \leq j \leq p$, which means that H_i are G -invariant.

The Hamiltonian vector fields X_{H_i}/G belong to $\mathcal{X}_{\mathcal{F}_H/G}$: Indeed, if $f \in \mathcal{F}_H/G$ then $\mathfrak{p}^*(f)$ is a first integral of H , implying $\{H_i, \mathfrak{p}^*(f)\} = 0$, or $\{H_i/G, f\} = 0$, where \mathfrak{p} denotes the projection $M \rightarrow M/G$.

To prove the integrability of X_H/G , it is sufficient to show that

$$(2.13) \quad \dim M/G \leq \text{ddim } \mathcal{F}_{H/G} + \text{ddim } (X_{H_1}/G, \dots, X_{H_q}/G)$$

But we denote by r the generic dimension of the intersection of a common level set of p independent first integrals of X_H with an orbit of G in M , then one can check that

$$p - \text{ddim } (X_{H_1}/G, \dots, X_{H_q}/G) = \text{ddim } \mathcal{X}_{\mathcal{F}_H} - \text{ddim } (X_{H_1}/G, \dots, X_{H_q}/G) = r$$

and

$$q - \text{ddim } \mathcal{F}_{H/G} = \text{ddim } \mathcal{F}_H - \text{ddim } \mathcal{F}_{H/G} \leq (d - k) - r$$

where $(d - k)$ is the dimension of a generic orbit of G in M . To prove the last inequality, notice that functions in $\mathcal{F}_{H/G}$ can be obtained from functions in \mathcal{F}_H by averaging with respect to the G -action. Also, G acts on the (separated) space of common level sets of the functions in \mathcal{F}_H , and isotropic groups of this G -action are of (generic) codimension $(d - k) - r$.

The above two formulas, together with $p + q = \dim M = \dim M/G + (d - k)$, implies Inequality (2.13) (it is in fact an equality). The proper case is straightforward. \square

2.5. Non-Hamiltonian reduced integrability.

One of the main differences between the non-Hamiltonian case and the Hamiltonian case is that reduced non-Hamiltonian integrability does not imply integrability. In fact, in the Hamiltonian case, we can lift Hamiltonian vector fields from M/G to M via the lifting of corresponding functions. In the non-Hamiltonian case, no such canonical lifting exists, therefore commuting vector fields on M/G do not provide commuting vector fields on M . For example, consider a vector field of the type $X = a_1 \partial/\partial x_1 + a_2 \partial/\partial x_2 + b(x_1, x_2) \partial/\partial x_3$ on the standard torus \mathbb{T}^3 with periodic coordinates (x_1, x_2, x_3) , where a_1 and a_2 are two incommensurable real numbers ($a_1/a_2 \notin \mathbb{Q}$), and $b(x_1, x_2)$ is a smooth function of two variables. Then clearly X is invariant under the \mathbb{T}^1 -action generated by $\partial/\partial x_3$, and the reduced system is integrable. On the other hand, for X to be integrable, we must be able to find a function $c(x_1, x_2)$ such that $[X, \partial/\partial x_1 + c(x_1, x_2) \partial/\partial x_3] = 0$. This last equation does not always have a solution (it is a small divisor problem, and depends on a_1/a_2 and the behavior of the coefficients of $b(x_1, x_2)$ in its Fourier expansion), i.e. there are choices of $a_1, a_2, b(x_1, x_2)$ for which the vector field X is not integrable.

However, non-Hamiltonian integrability still implies reduced integrability. Recall from Remark 2.9 that if a vector field X on a (generalized) manifold M is integrable, then under mild additional conditions we have $\text{ddim } \mathcal{X}_X + \text{ddim } \mathcal{F}_X = \dim M$, where \mathcal{F}_X is the set of all first integrals of X , and \mathcal{X}_X is the set of all vector fields which commute with X and preserve every function in \mathcal{F} .

Theorem 2.17. *Let X be a smooth non-Hamiltonian proper integrable system on a manifold M with the aid of $(\mathcal{F}_X, \mathcal{X}_X)$, i.e. $\text{ddim } \mathcal{X}_X + \text{ddim } \mathcal{F}_X = \dim M$, and G be a compact Lie group acting on M which preserves X . Then the reduced system on M/G is also proper integrable.*

Proof. Let \mathcal{X}_X^G denote the set of vector fields which belong to \mathcal{X}_X and which are invariant under the action of G . Note that the elements of \mathcal{X}_X^G can be obtained from the elements of \mathcal{X}_X by averaging with respect to the G -action.

A key ingredient of the proof is the fact $\text{ddim } \mathcal{X}_X^G = \text{ddim } \mathcal{X}_X$ (To see this fact, notice that near each regular invariant torus of the system there is an effective torus action (of the same dimension) which preserves the system, and this torus action must necessarily commute with the action of G . The generators of this torus action are linearly independent vector fields which belong to \mathcal{X}_X^G - in fact, they are defined locally near the union of G -orbits which by an invariant torus, but then we can extend them to global vector fields which lie in \mathcal{X}_X^G)

Therefore, we can project the pairwise commuting vector fields in \mathcal{X}_X^G from M to M/G to get pairwise commuting vector fields on M/G . To get the first integrals for the reduced system, we can also take the first integrals of X on M and average them with respect to the G -action to make them G -invariant. The rest of the proof of Theorem 2.17 is similar to that of Theorem 2.16. \square

3. TORUS ACTIONS AND LOCAL NORMAL FORMS

3.1. Toric characterization of Poincaré-Birkhoff normal form.

It is a simple well-known fact that every vector field near an equilibrium point admits a formal Poincaré-Birkhoff normal form (Birkhoff in the Hamiltonian case, and Poincaré-Dulac in the non-Hamiltonian case). What is also very simple but much less well-known is the fact that these normal forms are governed by torus actions. We will explain this fact here, following [94, 96].

Let X be a given analytic vector field in a neighborhood of 0 in \mathbb{K}^m , where $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , with $X(0) = 0$. When $\mathbb{K} = \mathbb{R}$, we may also view X as a holomorphic (i.e. complex analytic) vector field by complexifying it. Denote by

$$(3.1) \quad X = X^{(1)} + X^{(2)} + X^{(3)} + \dots$$

the Taylor expansion of X in some local system of coordinates, where $X^{(k)}$ is a homogeneous vector field of degree k for each $k \geq 1$.

In the Hamiltonian case, on a symplectic manifold, $X = X_H$, $m = 2n$, \mathbb{K}^{2n} has a standard symplectic structure, and $X^{(j)} = X_{H^{(j+1)}}$.

The algebra of linear vector fields on \mathbb{K}^m , under the standard Lie bracket, is nothing but the reductive algebra $gl(m, \mathbb{K}) = sl(m, \mathbb{K}) \oplus \mathbb{K}$. In particular, we have

$$(3.2) \quad X^{(1)} = X^s + X^{nil},$$

where X^s (resp., X^{nil}) denotes the semi-simple (resp., nilpotent) part of $X^{(1)}$. There is a complex linear system of coordinates (x_j) in \mathbb{C}^m which puts X^s into diagonal form:

$$(3.3) \quad X^s = \sum_{j=1}^m \gamma_j x_j \partial / \partial x_j,$$

where γ_j are complex coefficients, called **eigenvalues** of X (or $X^{(1)}$) at 0.

In the Hamiltonian case, $X^{(1)} \in sp(2n, \mathbb{K})$ which is a simple Lie algebra, and we also have the decomposition $X^{(1)} = X^s + X^{nil}$, which corresponds to the decomposition

$$(3.4) \quad H^{(2)} = H^s + H^{nil}$$

There is a complex canonical linear system of coordinates (x_j, y_j) in \mathbb{C}^{2n} in which H^s has diagonal form:

$$(3.5) \quad H^s = \sum_{j=1}^n \lambda_j x_j y_j,$$

where λ_j are complex coefficients, called **frequencies** of H (or $H^{(2)}$) at 0.

For each natural number $k \geq 1$, the vector field X^s acts linearly on the space of homogeneous vector fields of degree k by the Lie bracket, and the monomial vector fields are the eigenvectors of this action:

$$(3.6) \quad \left[\sum_{j=1}^m \gamma_j x_j \partial / \partial x_j, x_1^{b_1} x_2^{b_2} \dots x_n^{b_n} \partial / \partial x_l \right] = \left(\sum_{j=1}^n b_j \gamma_j - \gamma_l \right) x_1^{b_1} x_2^{b_2} \dots x_n^{b_n} \partial / \partial x_l.$$

When an equality of the type

$$(3.7) \quad \sum_{j=1}^m b_j \gamma_j - \gamma_l = 0$$

holds for some nonnegative integer m -tuple (b_j) with $\sum b_j \geq 2$, we will say that the monomial vector field $x_1^{b_1} x_2^{b_2} \dots x_m^{b_m} \partial / \partial x_l$ is a **resonant term**, and that the m -tuple $(b_1, \dots, b_l - 1, \dots, b_l)$ is a resonance relation for the eigenvalues (γ_i) . More precisely, a **resonance relation** for the n -tuple of eigenvalues (γ_j) of a vector field X is an m -tuple (c_j) of integers satisfying the relation $\sum c_j \gamma_j = 0$, such that $c_j \geq -1$, $\sum c_j \geq 1$, and at most one of the c_j may be negative.

In the Hamiltonian case, H^s acts linearly on the space of functions by the Poisson bracket. Resonant terms (i.e. generators of the kernel of this action) are monomials $\prod x_j^{a_j} y_j^{b_j}$ which satisfy the following resonance relation, with $c_j = a_j - b_j$:

$$(3.8) \quad \sum_{j=1}^m c_j \lambda_j = 0$$

Denote by \mathcal{R} the subset of \mathbb{Z}^m (or sublattice of \mathbb{Z}^n in the Hamiltonian case) consisting of all resonance relations (c_j) for a given vector field X . The number

$$(3.9) \quad r = \dim_{\mathbb{Z}}(\mathcal{R} \otimes \mathbb{Z})$$

is called the **degree of resonance** of X . Of course, the degree of resonance depends only on the eigenvalues of the linear part of X , and does not depend on the choice of local coordinates. If $r = 0$ then we say that the system is **nonresonant** at 0.

The vector field X is said to be in **Poincaré-Birkhoff normal form** if it commutes with the semisimple part of its linear part (see e.g. [14, 72]):

$$(3.10) \quad [X, X^s] = 0.$$

In the Hamiltonian case, the above equation can also be written as

$$(3.11) \quad \{H, H^s\} = 0.$$

The above equations mean that if X is in normal form then its nonlinear terms are resonant. A transformation of coordinates (which is symplectic in the Hamiltonian case) which puts X in Poincaré-Birkhoff normal form is called a **Poincaré-Birkhoff normalization**. It is a classical result of Poincaré, Dulac, and Birkhoff that any analytic vector field which vanishes at 0 admits a *formal* Poincaré-Birkhoff normalization (which does not converge in general).

Denote by $\mathcal{Q} \subset \mathbb{Z}^m$ the integral sublattice of \mathbb{Z}^m consisting of m -dimensional vectors $(\rho_j) \in \mathbb{Z}^m$ which satisfy the following properties :

$$(3.12) \quad \sum_{j=1}^m \rho_j c_j = 0 \quad \forall (c_j) \in \mathcal{R}, \text{ and } \rho_j = \rho_k \text{ if } \gamma_j = \gamma_k$$

(where \mathcal{R} is the set of resonance relations as before). In the Hamiltonian case, \mathcal{Q} is defined by

$$(3.13) \quad \sum_{j=1}^n \rho_j c_j = 0 \quad \forall (c_j) \in \mathcal{R}.$$

We will call the number

$$(3.14) \quad d = \dim_{\mathbb{Z}} \mathcal{Q}$$

the **toric degree** of X at 0. Of course, this number depends only on the eigenvalues of the linear part of X , and we have the following (in)equality : $r + d = n$ in the Hamiltonian case (where r is the degree of resonance), and $r + d \leq m$ in the non-Hamiltonian case.

Let $(\rho_j^1), \dots, (\rho_j^d)$ be a basis of \mathcal{Q} . For each $k = 1, \dots, d$ define the following diagonal linear vector field Z_k :

$$(3.15) \quad Z_k = \sum_{j=1}^m \rho_j^k x_j \partial / \partial x_j$$

in the non-Hamiltonian case, and $Z_k = X_{F^k}$ where

$$(3.16) \quad F^k = \sum_{j=1}^n \rho_j^k x_j y_j$$

in the Hamiltonian case.

The vector fields Z_1, \dots, Z_r have the following remarkable properties :

a) They commute pairwise and commute with X^s and X^{nil} , and they are linearly independent almost everywhere.

b) iZ_j is a periodic vector field of period 2π for each $j \leq r$ (here $i = \sqrt{-1}$). What does it mean is that if we write $iZ_j = \Re(iZ_j) + i\Im(iZ_j)$, then $\Re(iZ_j)$ is a periodic real vector field in $\mathbb{C}^n = \mathbb{R}^{2n}$ which preserves the complex structure.

c) Together, iZ_1, \dots, iZ_r generate an effective linear \mathbb{T}^r -action in \mathbb{C}^n (which preserves the symplectic structure in the Hamiltonian case), which preserves X^s and X^{nil} .

A simple calculation shows that X is in Poincaré-Birkhoff normal form, i.e. $[X, X^s] = 0$, if and only if we have

$$(3.17) \quad [X, Z_k] = 0 \quad \forall k = 1, \dots, r.$$

The above commutation relations mean that if X is in normal form, then it is preserved by the effective r -dimensional torus action generated by iZ_1, \dots, iZ_r . Conversely, if there is a torus action which preserves X , then because the torus is a compact group we can linearize this torus action (using Bochner's linearization theorem [9] in the non-Hamiltonian case, and the equivariant Darboux theorem in the Hamiltonian case, see e.g. [20, 39]), leading to a normalization of X . In other words, we have:

Theorem 3.1 ([94, 96]). *A holomorphic (Hamiltonian) vector field X in a neighborhood of 0 in \mathbb{C}^m (or \mathbb{C}^{2n} with a standard symplectic form) admits a locally holomorphic Poincaré-Birkhoff normalization if and only if it is preserved by an effective holomorphic (Hamiltonian) action of a real torus of dimension t , where t is the toric degree of $X^{(1)}$ as defined in (3.14), in a neighborhood of 0 in \mathbb{C}^m (or \mathbb{C}^{2n}), which has 0 as a fixed point and whose linear part at 0 has appropriate weights (given by the lattice \mathcal{Q} defined in (3.12,3.13), which depends only on the linear part $X^{(1)}$ of X).*

The above theorem is true in the formal category as well. But of course, any vector field admits a formal Poincaré-Birkhoff normalization, and a formal torus action.

3.2. Some simple consequences and generalizations.

Theorem 3.1 has many important implications. One of them is:

Proposition 3.2 ([94, 96]). *A real analytic vector field X (Hamiltonian or non-Hamiltonian) in the neighborhood of an equilibrium point admits a local real analytic Poincaré-Birkhoff normalization if and only if it admits a local holomorphic Poincaré-Birkhoff normalization when considered as a holomorphic vector field.*

The proof of the above proposition (see [94]) is based on the fact that the complex conjugation induces an involution on the torus action which governs the Poincaré-Birkhoff normalization.

If a dynamical system near an equilibrium point is invariant with respect to a compact group action which fixes the equilibrium point, then this compact group action commutes with the (formal) torus action of the Poincaré-Birkhoff normalization. Together, they form a bigger compact group action, whose linearization leads to a simultaneous Poincaré-Birkhoff normalization and linearization of the compact symmetry group, i.e. we can perform the Poincaré-Birkhoff normalization in an invariant way. This is a known result in dynamical systems, see e.g. [90], but the toric point of view gives a new simple proof of it. The case of equivariant vector fields is similar. For example, one can speak about Poincaré-Dulac normal forms for time-reversible vector fields, see e.g. [53].

Another situation where one can use the toric characterization is the case of isochore (i.e. volume preserving) vector fields. In this case, naturally, the normalization transformation is required to be volume-preserving. Both Theorem 3.1 and Proposition 3.2 remain valid in this case.

One can probably use the toric point of view to study normal forms of Hamiltonian vector field on *Poisson* manifolds as well. For example, let \mathfrak{g}^* be the dual of a semi-simple Lie algebra, equipped with the standard linear Poisson structure, and let $H : \mathfrak{g}^* \rightarrow \mathbb{K}$ be a regular function near the origin 0 of \mathfrak{g}^* . The corresponding Hamiltonian vector field X_H will vanish at 0, because the Poisson structure itself vanishes at 0. Applying Poincaré-Birkhoff normalization techniques, we can kill the “nonresonant terms” in H (with respect to the linear part of H , or $dH(0)$). The normalized Hamiltonian will be invariant under the coadjoint action of a subtorus of a Cartan torus of the (complexified) Lie group of \mathfrak{g} . In the “nonresonant” case, we have a Cartan torus action which preserves the system.

3.3. Convergent normalization for integrable systems.

Though every vector field near an equilibrium admits a formal Poincaré-Birkhoff normalization, the problem of finding a convergent (i.e. locally real analytic or holomorphic) normalization is much more difficult. The usual step by step killing of non-resonant terms leads to an infinite product of coordinate transformations, which may diverge in general, due to the presence of small divisors. Positive results about the convergence of this process are due to Poincaré, Siegel, Bruno and others mathematicians, under Diophantine conditions on the eigenvalues of the linear part of the system, see e.g. [14, 72].

However, when the vector field is analytically integrable (i.e. it is an real or complex analytic vector field, and the additional first integrals and commuting vector fields in question are also analytic), then we don’t need any Diophantine or nonresonance condition for the existence of a convergent Poincaré-Birkhoff normalization. More precisely, we have:

Theorem 3.3 ([94, 96]). *Let X be a local analytic (non-Hamiltonian, isochore, or Hamiltonian) vector field in $(\mathbb{K}^m, 0)$ (or in $(\mathbb{K}^{2n}, 0)$ with a standard symplectic structure), where $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , such that $X(0) = 0$. Then X admits a convergent Poincaré-Birkhoff normalization in a neighborhood of 0.*

Partial cases of the above theorem were obtained earlier by many authors, including Rüssmann [73] (the nondegenerate Hamiltonian case with 2 degrees of freedom), Vey [82, 83] (the nondegenerate Hamiltonian and isochore cases), Ito [42] (the nonresonant Hamiltonian case), Ito [44] and Kappeler et al. [48] (the Hamiltonian case with a simple resonance), Bruno and Walcher [15] (the non-Hamiltonian case with $m = 2$). These authors, except Vey who was more geometric, relied on long and heavy analytical estimates to show the convergence of an infinite normalizing coordinate transformation process. On the other hand, the proof of Theorem 3.3 in [94, 96] is based on the toric point of view and is relatively short.

Following [94], we will give here a sketch of the proof of the above theorem in the Liouville-integrable case. The other cases are similar, and of course the theorem is valid for Hamiltonian vector fields which are integrable in generalized Liouville sense as well. According to Proposition 3.2, it is enough to show the existence of a holomorphic normalization. We will do it by finding local Hamiltonian \mathbb{T}^1 -actions which preserve the moment map of an analytically completely integrable system. The Hamiltonian function generating such an action is an **action function**. If we find $(n - q)$ such \mathbb{T}^1 -actions, then they will automatically commute and give rise to a Hamiltonian \mathbb{T}^{n-q} -action.

To find an action function, we will use the Mineur-Arnold formula $P = \int_{\Gamma} \beta$, where P denotes an action function, β denotes a primitive 1-form (i.e. $\omega = d\beta$ is the symplectic form), and Γ denotes an 1-cycle (closed curve) lying on a level set of the moment map. To show the existence of such 1-cycles Γ , we will use an approximation method, based on the existence of a formal Birkhoff normalization.

Denote by $\mathbf{G} = (G_1 = H, G_2, \dots, G_n) : (\mathbb{C}^{2n}, 0) \rightarrow (\mathbb{C}^n, 0)$ the holomorphic momentum map germ of a given complex analytic Liouville-integrable Hamiltonian system. Let $\epsilon_0 > 0$ be a small positive number such that \mathbf{G} is defined in the ball $\{z = (x_j, y_j) \in \mathbb{C}^{2n}, |z| < \epsilon_0\}$. We will restrict our attention to what happens inside this ball. As in Subsection 3.1, we may assume that in the symplectic coordinate system $z = (x_j, y_j)$ we have

$$(3.18) \quad H = G_1 = H^s + H^n + H^{(3)} + H^{(4)} + \dots$$

with

$$(3.19) \quad H_s = \sum_{k=1}^{n-q} \alpha_k F^k, \quad F^k = \sum_{j=1}^n \rho_j^k x_j y_j,$$

with no resonance relations among $\alpha_1, \dots, \alpha_{n-q}$. We will fix this coordinate system $z = (x_j, y_j)$, and all functions will be written in this coordinate system.

The real and imaginary parts of the Hamiltonian vector fields of G_1, \dots, G_n are in involution and their infinitesimal \mathbb{C}^n -action defines an **associated singular Lagrangian fibration** in the ball $\{z = (x_j, y_j) \in \mathbb{C}^{2n}, |z| < \epsilon_0\}$. For each z we will denote the fiber which contains z by M_z . If z is a point such that $\mathbf{G}(z)$ is a regular value for the momentum map, then M_z is a connected component of $\mathbf{G}^{-1}(\mathbf{G}(z))$.

Denote by

$$(3.20) \quad S = \{z \in \mathbb{C}^{2n}, |z| < \epsilon_0, dG_1 \wedge dG_2 \wedge \dots \wedge dG_n(z) = 0\}$$

the singular locus of the moment map, which is also the set of singular points of the associated singular foliation. What we need to know about S is that it is analytic and of codimension at least 1, though for generic integrable systems S is in fact of codimension 2. In particular, we have the following Lojasiewicz inequality [59]: there exist a positive number N and a positive constant C such that

$$(3.21) \quad |dG_1 \wedge \dots \wedge dG_n(z)| > C(d(z, S))^N$$

for any z with $|z| < \epsilon_0$, where the norm applied to $dG_1 \wedge \dots \wedge dG_n(z)$ is some norm in the space of n -vectors, and $d(z, S)$ is the distance from z to S with respect to the Euclidean metric. In the above inequality, if we change the coordinate system, then only ϵ_0 and C have to be changed, N (the Lojasiewicz exponent) remains the same.

We will choose an infinite decreasing series of small numbers ϵ_m ($m = 1, 2, \dots$), as small as needed, with $\lim_{m \rightarrow \infty} \epsilon_m = 0$, and define the following open subsets U_m of \mathbb{C}^{2n} :

$$(3.22) \quad U_m = \{z \in \mathbb{C}^{2n}, |z| < \epsilon_m, d(z, S) > |z|^m\}$$

We will also choose two infinite increasing series of natural numbers a_m and b_m ($m = 1, 2, \dots$), as large as needed, with $\lim_{m \rightarrow \infty} a_m = \lim_{m \rightarrow \infty} b_m = \infty$. It follows from Birkhoff's formal normalization that there is a series of local holomorphic

symplectic coordinate transformations Φ_m , $m \in \mathbb{N}$, such that the following two conditions are satisfied :

a) The differential of Φ_m at 0 is identity for each m , and for any two numbers m, m' with $m' > m$ we have

$$(3.23) \quad \Phi_{m'}(z) = \Phi_m(z) + O(|z|^{a_m}).$$

In particular, there is a formal limit $\Phi_\infty = \lim_{m \rightarrow \infty} \Phi_m$.

b) The moment map is normalized up to order b_m by Φ_m . More precisely, the functions G_j can be written as

$$(3.24) \quad G_j(z) = G_{(m)j}(z) + O(|z|^{b_m}), \quad j = 1, \dots, n,$$

with $G_{(m)j}$ such that

$$(3.25) \quad \{G_{(m)j}, F_{(m)}^k\} = 0 \quad \forall j = 1, \dots, n, \quad k = 1, \dots, n - q.$$

Here the functions $F_{(m)}^k$ are quadratic functions

$$(3.26) \quad F_{(m)}^k(x, y) = \sum_{j=1}^n \rho_j^k x_{(m)j} y_{(m)j}$$

in local symplectic coordinates

$$(3.27) \quad (x_{(m)}, y_{(m)}) = \Phi_m(x, y).$$

Notice that $F_{(m)}^k$ has the same form as F^k , but with respect to a different coordinate system. When considered in the original coordinate system (x, y) , $F_{(m)}^k$ is a different function than F^k , but the quadratic part of $F_{(m)}^k$ is F^k .

Denote by $\Gamma_m^k(z)$ the orbit of the real part of the periodic Hamiltonian vector field $X_{iF_{(m)}^k}$ which goes through z . Then for any $z' \in \Gamma_m^k(z)$ we have $G_{(m)j}(z') = G_{(m)j}(z)$ and $|z'| \simeq |z|$, therefore

$$(3.28) \quad |\mathbf{G}(z') - \mathbf{G}(z)| = O(|z'|^{b_m}).$$

(Note that we can choose the numbers a_m and b_m first, then choose the radii ϵ_m of small open subsets to make them sufficiently small with respect to a_m and b_m , so that the equivalence $O(|z'|^{b_m}) \simeq O(|z|^{b_m})$ makes sense). It follows from the definition of U_m and Lojasiewicz inequalities that we also have

$$(3.29) \quad |dG_1(z') \wedge \dots \wedge dG_n(z')| > d(z, S)^N > |z|^{mN}$$

for any $z \in U_m$ and $z' \in \Gamma_m^k(z)$, provided that b_m is sufficiently large and ϵ_m is sufficiently small. Assuming that $b_m \gg mN$, we can project the curve $\Gamma_m^k(z)$ on M_z in a unique natural way up to homotopy. Denote the image of the projection by $\tilde{\Gamma}_m^k(z)$, and define the following function P_m^k on U_m :

$$(3.30) \quad P_m^k(z) = \oint_{\tilde{\Gamma}_m^k(z)} \sum_{j=1}^n x_j dy_j .$$

One then checks that P_m^k is a uniformly bounded (say by 1) holomorphic first integral of the system on U_m , and moreover P_m^k coincides with $P_{m'}^k$ on $U_m \cap U_{m'}$ for

any m, m' , and hence we have a holomorphic first integral P^k on $U = \bigcup_{m=1}^{\infty} U_m$. The following lemma 3.4 about holomorphic extension says that P^k can be extended to a holomorphic first integral of the system in a neighborhood of 0. It is easy to see that P^k is an action function (because $P^k = \lim_{m \rightarrow \infty} \sqrt{-1} F_{(m)}^k$), i.e. its corresponding Hamiltonian flow is periodic of period 2π . Since $k = 1, \dots, n - q$, we have $n - q$ action functions, whose flows commute and generate the required Hamiltonian \mathbb{T}^{n-q} -action which preserves the system. \square

The following lemma on holomorphic extension, which is interesting in its own right, implies that the action functions P^k constructed above can be extended holomorphically to a neighborhood of 0.

Lemma 3.4. *Let $U = \bigcup_{m=1}^{\infty} U_m$, with $U_m = \{x \in \mathbb{C}^n, |x| < \epsilon_m, d(x, S) > |x|^m\}$, where ϵ_m is an arbitrary series of positive numbers and S is a local proper complex analytic subset of \mathbb{C}^n ($\text{codim}_{\mathbb{C}} S \geq 1$). Then any bounded holomorphic function on U has a holomorphic extension to a neighborhood of 0 in \mathbb{C}^n .*

See [94] for the proof of Lemma 3.4. It is a straightforward proof in the case S is non-singular or is a normal crossing, and makes use of a desingularization of S in the general case. \square

3.4. Torus action near a compact singular orbit.

Consider a real analytic integrable vector field X on a real analytic manifold M^m of dimension $m = p + q$, with the aid of a p -tuple $\mathbf{X} = (X_1, \dots, X_p)$ of commuting analytic vector fields and a q -tuple $\mathbf{F} = (F_1, \dots, F_q)$ of analytic common first integrals: $[X, X_i] = [X_i, X_j] = 0$, $X(F_j) = X_i(F_j) = 0 \forall i, j$. In the Hamiltonian case, when there is an analytic Poisson structure on M^m , we suppose that the system is integrable in generalized Liouville sense, i.e. the vector fields X, X_1, \dots, X_p are Hamiltonian.

The commuting vector fields X_1, \dots, X_p generate an infinitesimal \mathbb{R}^p -action on M – as usual, its orbits will be called orbits of the system. The map $\mathbf{F} : M^m \rightarrow \mathbb{R}^q$ is constant on the orbits of the system. Let $O \subset M^m$ be a singular orbit of dimension r of the system, $0 \leq r < p$. We suppose that O is a compact submanifold of M^m (or more precisely, of the interior of M^m if M^m has boundary). Then O is a torus of dimension r . Denote by N the connected component of $\mathbf{F}^{-1}(\mathbf{F}(O))$ which contains O . A natural question arises: does there exist a \mathbb{T}^r -action in a neighborhood of O or N , which preserves the system and is transitive on O ?

The above question has been answered positively in [98], under a mild condition called the *finite type condition*. To formulate this condition, denote by $M_{\mathbb{C}}$ a small open complexification of M^m on which the complexification $\mathbf{X}_{\mathbb{C}}, \mathbf{F}_{\mathbb{C}}$ of \mathbf{X} and \mathbf{F} exists. Denote by $N_{\mathbb{C}}$ a connected component of $\mathbf{F}_{\mathbb{C}}^{-1}(\mathbf{F}_{\mathbb{C}}(O))$ which contains N .

Definition 3.5. With the above notations, the singular orbit O is called of **finite type** if there is only a finite number of orbits of the infinitesimal action of \mathbb{C}^p in $N_{\mathbb{C}}$, and $N_{\mathbb{C}}$ contains a regular point of the map \mathbf{F} .

For example, all nondegenerate singular orbits are of finite type (see Section 4). We conjecture that every singular orbit of an algebraically integrable system is of finite type.

Theorem 3.6 ([98]). *With the above notations, if O is a compact finite type singular orbit of dimension r , then there is a real analytic torus action of \mathbb{T}^r in a neighborhood of O which preserves the integrable system (\mathbf{X}, \mathbf{F}) and which is transitive on O . If moreover N is compact, then this torus action exists in a neighborhood of N . In the Hamiltonian case this torus action also preserves the Poisson structure.*

Notice that Theorem 3.6, together with Theorem 3.3 and the toric characterization of Poincaré-Birkhoff normalization, provides an analytic Poincaré-Birkhoff normal form in the neighborhood a singular invariant torus of an integrable system.

Denote by \mathcal{A}_O the local automorphism group of the integrable system (\mathbf{X}, \mathbf{F}) at O , i.e. the group of germs of local analytic automorphisms of (\mathbf{X}, \mathbf{F}) in vicinity of O (which preserve the Poisson structure in the Hamiltonian case). Denote by \mathcal{A}_O^0 the subgroup of \mathcal{A}_O consisting of elements of the type g_Z^1 , where Z is a analytic vector field in a neighborhood of O which preserves the system and g_Z^1 is the time-1 flow of Z . The torus in the previous theorem is of course a Abelian subgroup of \mathcal{A}_O^0 . Actually, the automorphism group \mathcal{A}_O itself is essentially Abelian in the finite type case:

Theorem 3.7 ([98]). *If O is a compact finite type singular orbit as above, then \mathcal{A}_O^0 is an Abelian normal subgroup of \mathcal{A}_O , and $\mathcal{A}_O/\mathcal{A}_O^0$ is a finite group.*

The above two theorems are very closely related: their proofs are almost the same. Let us indicate here the main ingredients of the proof of Theorem 3.6:

For simplicity, we will assume that $r = 1$, i.e. O is a circle (the case $r > 1$ is absolutely similar). Since O is of finite type, there is a regular complex orbit Q in $N_{\mathbb{C}}$ of dimension p whose closure contains O . Q is a flat affine manifold (the affine structure is given by the \mathbb{C}^p -action, so we can talk about geodesics on Q . If we can find a closed geodesic γ_Q on Q , then it is a periodic orbit of period 1 of a vector field of the type $\sum a_j X_j$ on Q (with a_j being constants) on Q . Since the points of Q are regular for the map \mathbf{F} , using implicit function theorem, we can construct a vector field of the type $\sum a_j X_j$, with a_j now being holomorphic functions which are functionally dependent on \mathbf{F} (so that this vector fields preserves the system), and which is periodic of period 1 near γ_Q . With some luck, we will be able to extend this vector field holomorphically to a vector field in a neighborhood of O so that O becomes a periodic orbit of it, and we are almost done: if the vector field is not real-analytic, then its image under a complex involution will be another periodic vector field which preserves the system; the two vector fields commute (because the system is integrable) and we can fabricate from them a real-analytic periodic vector field, i.e. a real-analytic \mathbb{T}^1 -action in a neighborhood of O , for which O is a periodic orbit.

The main difficulty lies in finding the closed geodesic γ_Q (which satisfies some additional conditions). We will do it inductively: let $O_1 = O_{\mathbb{C}}$ ($O \subset O_{\mathbb{C}}$), $O_2, \dots, O_k = Q$ be a maximal chain of complex orbits of the system in $N_{\mathbb{C}}$ such that O_i lies in the closure of O_{i+1} and $O_i \neq O_{i+1}$. Then on each O_i , we will find a closed geodesic γ_i , such that each γ_{i+1} is homotopic to a multiple of γ_i in $O_i \cup O_{i+1}$, starting with $\gamma_1 = O$. We will show how to go from $O = \gamma_1$ to γ_2 (the other steps are similar). Without loss of generality, we may assume that O is a closed orbit for X_1 . Take a small section D to O in M , and consider the Poincare map ϕ of X_1 on D . Let $Y = O_2 \cap D_{\mathbb{C}}$. Then Y is a affine manifold (whose affine structure is projected from

O_2 by X_1). Let y be a point in Y . We want to connect y to $\phi(y)$ by a geodesic in Y . If we can do it, then the sum of this geodesic segment with the orbit of X_1 going from y to $\phi(Y)$ can be modified into a closed geodesic γ_2 on O_2 . Unfortunately, in general, we cannot connect y to $\phi(y)$ by a geodesic in Y , because Y is not “convex”. But a lemma says that Y can be cut into a finite number of convex pieces, and as a consequence y can be connected geodesically to $\phi^N(y)$ for some power ϕ^N (N -time iteration) of ϕ . See [98] for the details. \square

Theorem 3.6 reduces the study of the behavior of integrable systems near compact singular orbits to the study of fixed points with a finite Abelian group of symmetry (this group arises from the fact that the torus action is not free in general, only locally free). For example, as was shown in [93], the study of *corank-1* singularities of Liouville-integrable systems is reduced to the study of families of functions on a 2-dimensional symplectic disk which are invariant under the rotation action of a finite cyclic group \mathbb{Z}/\mathbb{Z}_k , where one can apply the theory of singularities of functions with an Abelian symmetry developed by Wassermann [84] and other people. A (partial) classification up to diffeomorphisms of corank-1 degenerate singularities was obtained by Kalashnikov [47] (see also [93, 60]), and symplectic invariants were obtained by Colin de Verdière [17].

4. NONDEGENERATE SINGULARITIES

In this section, we will consider only smooth Liouville-integrable Hamiltonian systems, though many ideas and results can probably be extended to other kinds of integrable systems.

4.1. Nondegenerate singular points.

Consider the momentum map $\mathbf{F} = (F_1, \dots, F_n) : (M^{2n}, \omega) \rightarrow \mathbb{R}^n$ of a smooth integrable Hamiltonian system on a symplectic manifold (M^{2n}, ω) . In this Section, we will forget about the original Hamiltonian function, and study the momentum map instead.

For a point $z \in M$, denote $\text{rank } z = \text{rank } d\mathbf{F}(z)$, where $d\mathbf{F}$ denotes the differential of \mathbf{F} . This number is equal to the dimension of the orbit of the system (i.e. the infinitesimal Poisson \mathbb{R}^n -action generated by X_{F_1}, \dots, X_{F_n}) which goes through z . If $\text{rank } z < n$ then z is called a **singular point**. If $\text{rank } z = 0$ then z is a **fixed point** of the system.

If z is a fixed point, then the quadratic parts $F_1^{(2)}, \dots, F_n^{(2)}$ of the components F_1, \dots, F_n of the momentum map at z are Poisson-commuting and they form an Abelian subalgebra, A_z , of the Lie algebra $Q(2n, \mathbb{R})$ of homogeneous quadratic functions of $2n$ variables under the standard Poisson bracket. Observe that the algebra $Q(2n, \mathbb{R})$ is isomorphic to the symplectic algebra $sp(2n, \mathbb{R})$.

A fixed point z will be called **nondegenerate** if A_z is a Cartan subalgebra of $Q(2n, \mathbb{R})$. In this case, according to Williamson [87], there is a triple of nonnegative integers (k_e, k_h, k_f) such that $k_e + k_h + 2k_f = n$, and a canonical coordinate system (x_i, y_i) in \mathbb{R}^{2n} , such that A_z is spanned by the following quadratic functions

h_1, \dots, h_n :

$$(4.1) \quad \begin{aligned} h_i &= x_i^2 + y_i^2 \text{ for } 1 \leq i \leq k_e ; \\ h_i &= x_i y_i \text{ for } k_e + 1 \leq i \leq k_e + k_h ; \\ h_i &= x_i y_{i+1} - x_{i+1} y_i \text{ and} \\ h_{i+1} &= x_i y_i + x_{i+1} y_{i+1} \text{ for } i = k_e + k_h + 2j - 1, 1 \leq j \leq k_f . \end{aligned}$$

The triple (k_e, k_h, k_f) is called the **Williamson type** of (the system at) z . k_e is the number of **elliptic** components (and h_1, \dots, h_{k_e} are elliptic components), k_h is the number of **hyperbolic** components, and k_f is the number of **focus-focus** components. If $k_h = k_f = 0$ then z is called an **elliptic singular point**.

The local structure of nondegenerate singular points is given by the following theorem of Eliasson.

Theorem 4.1 (Eliasson [29, 30]). *If z is a nondegenerate fixed point of a smooth Liouville-integrable Hamiltonian system then there is a smooth Birkhoff normalization. In other words, the singular Lagrangian foliation given by the momentum map \mathbf{F} in a neighborhood of z is locally smoothly symplectomorphic to the “linear” singular Lagrangian fibration given by the quadratic map $(h_1, \dots, h_n) : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ with the standard symplectic structure on \mathbb{R}^{2n} .*

The elliptic case of the above theorem is also obtained independently by Dufour and Molino [25]. The case one degree of freedom is due to Colin de Verdière and Vey [18]. The analytic case of the above theorem is due to Vey [82], and is superseded by Theorem 3.3. There is also a semiclassical version of Eliasson’s theorem (quantum Birkhoff normal form), which is due to Vu Ngoc San [75].

The proof of Eliasson’s theorem [29, 30] is quite long and highly technical: The first step is to use division lemmas in singularity theory to show that the local singular fibration given by the momentum map is diffeomorphic (without the symplectic structure) to the linear model. Then one uses a combination of averaging, Moser’s path method, and technics similar to the ones used in the proof of Sternberg’s smooth linearization theorem for vector fields, to show that the symplectic form can also be normalized smoothly. In fact, Eliasson’s proof of his theorem is not quite complete, except for the elliptic case, because it lacks some details which were difficult to work out, see [65].

A direct consequence of Eliasson’s theorem is that, near a nondegenerate fixed point of Williamson type (k_e, k_h, k_f) , there is a local smooth Hamiltonian $\mathbb{T}^{k_e+k_f}$ -action which preserves the system: each elliptic or focus-focus component provides one \mathbb{T}^1 -action. In the analytic case, Birkhoff normalization gives us a \mathbb{T}^n -action, but it acts in the complex space, and in the real space we only see a $\mathbb{T}^{k_e+k_f}$ -action.

4.2. Nondegenerate singular orbits.

Let $x \in M$ be a singular point of rank $x = m \geq 0$. We may assume without loss of generality that $dF_1 \wedge \dots \wedge dF_m(x) \neq 0$, and a local symplectic reduction near x with respect to the local free \mathbb{R}^m -action generated by the Hamiltonian vector fields X_{F_1}, \dots, X_{F_m} will give us an m -dimensional family of local integrable Hamiltonian systems with $n - m$ degrees of freedom. Under this reduction, x will be mapped to a fixed point in the reduced system, and if this fixed point is nondegenerate according to the above definition, then x is called a **nondegenerate singular**

point of rank m and corank $(n - m)$. In this case, we can speak about the Williamson type (k_e, k_h, k_f) of x , and we have $k_e + k_h + 2k_f = m$.

A **nondegenerate singular orbit** of the system is an orbit (of the infinitesimal Poisson \mathbb{R}^n -action) which goes through a nondegenerate singular point. Since all points on a singular orbit have the same Williamson type, we can speak about the Williamson type and the corank of a nondegenerate singular orbit. We have the following generalization of Theorem 4.1 to the case of compact nondegenerate singular orbits:

Theorem 4.2 (Miranda–Zung [66]). *If O is a compact nondegenerate singular orbit of a smooth Liouville-integrable Hamiltonian system, then the singular Lagrangian fibration given by the momentum map in a neighborhood of O is smoothly symplectomorphic to a linear model. Moreover, if the system is invariant under a symplectic action of a compact Lie group G in a neighborhood of O , then the above smooth symplectomorphism to the linear model can be chosen to be G -equivariant.*

The linear model in the above theorem can be constructed as follows: Denote by (p_1, \dots, p_m) a linear coordinate system of a small ball D^m of dimension m , $(q_1 \pmod{1}, \dots, q_m \pmod{1})$ a standard periodic coordinate system of the torus \mathbb{T}^m , and $(x_1, y_1, \dots, x_{n-m}, y_{n-m})$ a linear coordinate system of a small ball $D^{2(n-m)}$ of dimension $2(n - m)$. Consider the manifold

$$(4.2) \quad V = D^m \times \mathbb{T}^m \times D^{2(n-m)}$$

with the standard symplectic form $\sum dp_i \wedge dq_i + \sum dx_j \wedge dy_j$, and the following momentum map: $(\mathbf{p}, \mathbf{h}) = (p_1, \dots, p_m, h_1, \dots, h_{n-m}) : V \rightarrow \mathbb{R}^n$, where (h_1, \dots, h_{n-m}) are quadratic functions given by Equation (4.1). A symplectic group action on V which preserves the above momentum map is called *linear* if it on the product $V = D^m \times \mathbb{T}^m \times D^{2(n-m)}$ componentwise, the action on D^m is trivial, the action on \mathbb{T}^m is by translations with respect to the coordinate system (q_1, \dots, q_m) , and the action on $D^{2(n-m)}$ is linear.

Let Γ be a finite group with a free linear symplectic action $\rho(\Gamma)$ on V which preserves the momentum map. Then we can form the quotient integrable system with the momentum map

$$(4.3) \quad (\mathbf{p}, \mathbf{h}) = (p_1, \dots, p_m, h_1, \dots, h_{n-m}) : V/\Gamma \rightarrow \mathbb{R}^n .$$

The set $\{p_i = x_i = y_i = 0\} \subset V/\Gamma$ is a compact orbit of Williamson type (k_e, k_f, k_h) of the above system. The above system on V/Γ is called the **linear model** of Williamson type (k_e, k_f, k_h) and twisting group Γ , or more precisely, twisting action $\rho(\Gamma)$. (It is called a direct model if Γ is trivial, and a twisted model if Γ is nontrivial). A symplectic action of a compact group G on V/Γ which preserves the momentum map $(p_1, \dots, p_m, h_1, \dots, h_{n-m})$ is called linear if it comes from a linear symplectic action of G on V which commutes with the action of Γ .

The case with G trivial and $n = 2, k_h = 1, k_e = k_f = 0$ of Theorem 4.2 is due to Colin de Verdière and Vu Ngoc San [19], and independently Currás-Bosch and Miranda [21]. A direct consequence of Theorem 4.2 is that the group of local smooth symplectic automorphisms of a smooth Liouville-integrable system near a compact nondegenerate singular orbit is Abelian, see [66].

4.3. Nondegenerate singular fibers.

In this subsection, we will assume that the momentum map $\mathbf{F} : M^{2n} \rightarrow \mathbb{R}^n$ is proper. A singular connected component of a level set of the momentum map will be called a **singular fiber** of the system. A singular fiber may contain one orbit (e.g. in the elliptic nondegenerate case), or many orbits, some of them singular and some of them regular. A singular fiber N_c is called **nondegenerate** if any point $z \in N_c$ is either regular or nondegenerate singular. The nondegeneracy is an open condition: if a singular fiber is nondegenerate then nearby singular fibers are also nondegenerate.

By a **singularity** of a Liouville-integrable system, we mean the germ of the system near a singular fiber, together with the symplectic form and the Lagrangian fibration. We will denote a singularity by $(\mathcal{U}(N_c), \omega, \mathcal{L})$, where $\mathcal{U}(N_c)$ denotes a small “tubular” neighborhood of N_c , and \mathcal{L} denotes the Lagrangian fibration. If N_c is nondegenerate then $(\mathcal{U}(N_c), \omega, \mathcal{L})$ is also called nondegenerate.

A simple lemma [92] says that if N_c is a nondegenerate singular fiber, then all singular points of maximal corank in N_c have the same Williamson type. We define the rank and the Williamson type of a nondegenerate singularity $(\mathcal{U}(N_c), \omega, \mathcal{L})$ to be the rank and the Williamson type of a singular point of maximal corank in N_c .

The following theorem may be viewed as the generalization of Liouville–Mineur–Arnold theorem to the case of nondegenerate singular fibers:

Theorem 4.3 ([92]). *Let $(\mathcal{U}(N_c), \omega, \mathcal{L})$ be a nondegenerate smooth singularity of rank m and Williamson type (k_e, k_h, k_f) of a Liouville-integrable system with a proper momentum map. Then we have:*

- a) *There is effective Hamiltonian $\mathbb{T}^{m+k_e+k_f}$ -action in $(\mathcal{U}(N_c), \omega, \mathcal{L})$ which preserves the system. The dimension $m + k_e + k_f$ is maximal possible. There is a locally free \mathbb{T}^m -subaction of this action.*
- b) *There is a partial action-angle coordinate system.*
- c) *Under a mild additional condition, $(\mathcal{U}(N_c), \mathcal{L})$ is topologically equivalent to an almost direct product of simplest (corank 1 elliptic or hyperbolic and corank 2 focus-focus) singularities.*

Assertion b) of the above theorem means that we can write $(\mathcal{U}(N_c), \omega)$ as $(D^m \times \mathbb{T}^m \times P^{2k})/\Gamma$ with

$$(4.4) \quad \omega = \sum_1^m dp_i \wedge dq_i + \omega_1$$

where ω_1 is a symplectic form on P^{2k} , the finite group Γ acts on the product component-wise, its action is linear on \mathbb{T}^m , and the momentum map \mathbf{F} does not depend on the variables q_1, \dots, q_m .

The additional condition in Assertion c) prohibits the bifurcation diagram (i.e. the set of singular values of the momentum map) from having “pathologies”, see [92], and it’s satisfied for all nondegenerate singularities of physical integrable systems met in practice. The almost direct product means a product of the type

$$(4.5) \quad (\mathcal{T}^{2m} \times \mathcal{E}_1^2 \times \dots \times \mathcal{E}_{k_e}^2 \times \mathcal{H}_1^2 \times \dots \times \mathcal{H}_{k_h}^2 \times \mathcal{F}_1^4 \times \dots \times \mathcal{F}_{k_f}^4)/\Gamma$$

where \mathcal{T}^{2m} is the germ of $(D^m \times \mathbb{T}^m, \sum_1^m dp_i \wedge dq_i)$ with the standard Lagrangian torus fibration; $\mathcal{E}_i^2, \mathcal{F}_i^2$ and \mathcal{H}_i^4 are elliptic, hyperbolic and focus-focus singularities of integrable systems on symplectic manifolds of dimension 2, 2 and 4 respectively; the finite group Γ acts freely and component-wise. Remark that, in general, a nondegenerate singularity is only topologically equivalent, but not symplectically equivalent, to an almost direct product singularity.

The above almost direct product may remind one of the decomposition of algebraic reductive groups into almost direct products of simple groups and tori: though the two objects are completely different, there are some common ideas behind them, namely infinitesimal direct decomposition, and twisting by a finite group.

4.4. Focus-focus singularities.

The singularities $\mathcal{E}_i^2, \mathcal{H}_i^2, \mathcal{F}_i^4$ in (4.5) may be called **elementary** nondegenerate singularities; they are characterized by the fact that $k_e + k_h + k_f = 1$ and $\text{rank} = 0$. Among them, elliptic singularities \mathcal{E}_i^2 are very simple: each elementary elliptic singularity is isomorphic to a standard linear model (a harmonic oscillator). Elementary hyperbolic singularities \mathcal{H}_i^2 are also relatively simple because they are given by hyperbolic singular level sets of Morse functions on 2-dimensional symplectic surfaces. On the other hand, *focus-focus* singularities \mathcal{F}_i^4 live in 4-dimensional symplectic manifolds, so their topological structure is somewhat more interesting. Let us mention here some results about the structure of these 4-dimensional focus-focus singularities, see [92, 95] and references therein for more details.

One of the most important facts about focus-focus singularities is the existence of a \mathbb{T}^1 -action (this is a special case of Assertion a) of Theorem 4.3); many other important properties are consequences of this \mathbb{T}^1 -action. In fact, in many integrable systems with a focus-focus singularity, e.g. the spherical pendulum and the Lagrangian top, this \mathbb{T}^1 -action is the obvious rotational symmetry, though in some systems, e.g. the Manakov integrable system on $so(4)$, this local \mathbb{T}^1 -action is “hidden”. Dynamically speaking, a focus-focus point is roughly an unstable equilibrium point with a \mathbb{T}^1 -symmetry.

Each focus-focus singularity has only one singular fiber: the focus-focus fiber, which is homeomorphic to a **pinched torus** (take a torus, and ℓ parallel homotopically non-trivial simple closed curves on it, $\ell \geq 1$, then collapse each of these curves into one point). This fact was known to Lerman and Umanskij [54, 55].

From the topological point of view, we have a singular torus fibration in a four-dimensional manifold with one singular fiber. These torus fibrations have been studied by Matsumoto and other people, see e.g. [61] and references therein, and of course the case with a singular fiber of focus-focus type is included in their topological classification. In particular, the number of pinches ℓ is the only topological invariant. The monodromy of the torus fibration (over a punched 2-dimensional disk) around the focus-focus fiber is given by the matrix $\begin{pmatrix} 1 & \ell \\ 0 & 1 \end{pmatrix}$. By the way, the case with $\ell > 1$ is topologically an ℓ -sheet covering of the case with $\ell = 1$, and a concrete example with $\ell = 1$ is the unstable equilibrium of the usual spherical pendulum. This phenomenon of *nontrivial monodromy* (of the foliation by Liouville tori) was first observed by Duistermaat and Cushman [26], and then by some other people for various concrete integrable systems. Now we have many different

ways to look at this monodromy: from the purely topological point of view (using Matsumoto's theory [61]), from the point of view of Picard-Lefschetz theory (see Audin [6] and references therein), or as a consequence of Duistermaat-Heckman formula with respect to the above-mentioned \mathbb{T}^1 -action (see [95]). Quantization of focus-focus singularities leads to *quantum* monodromy, see Vu Ngoc San [74] and Section 5.

Similar results, including the existence of a \mathbb{T}^1 -action, for focus-focus singularities of *non-Hamiltonian* integrable systems, have been obtained by Cushman and Duistermaat [22], see also [95].

5. GLOBAL ASPECTS OF LOCAL TORUS ACTIONS

5.1. Sheaf of local \mathbb{T}^1 -actions.

Consider a smooth proper integrable system on a manifold M with a given p -tuple of commuting vector fields $\mathbf{X} = (X_1, \dots, X_p)$ and q -tuple of common first integrals $\mathbf{F} = (F_1, \dots, F_q)$.

We will call the space of connected components of the level sets of the map \mathbf{F} the **base space** of the integrable system, and denote it by \mathcal{B} . Since the system is proper, the space \mathcal{B} with the induced topology from M is a Hausdorff space. We will denote by $\mathbf{P} : M \rightarrow \mathcal{B}$ the projection map from M to \mathcal{B} .

For each open set U of \mathcal{B} , denote by $\mathcal{R}(U)$ the set of all \mathbb{T}^1 -actions in $\mathbf{P}^{-1}(U)$ which preserve the integrable system (\mathbf{X}, \mathbf{F}) (in the Hamiltonian case, due to generalized Liouville-Mineur-Arnold theorem, elements of $\mathcal{R}(U)$ will automatically preserve the Poisson structure). $\mathcal{R}(U)$ is an Abelian group: if two elements ρ_1, ρ_2 of $\mathcal{R}(U)$ are generated by two periodic vector fields Y_1, Y_2 respectively, then Y_1 will automatically commute with Y_2 , and the sum $Y_1 + Y_2$ generates another \mathbb{T}^1 -action which can be called the sum of ρ_1 and ρ_2 . Actually, $\mathcal{R}(U)$ is a free Abelian group, and its dimension can vary from 0 to p (the dimension of a regular invariant torus of the system), depending on U and on the system. If U is a small disk in the regular region of \mathcal{B} then $\dim_{\mathbb{Z}} \mathcal{R}(U) = p$.

The association $U \mapsto \mathcal{R}(U)$ forms a free Abelian sheaf \mathcal{R} over \mathcal{B} , which we will call the **toric monodromy sheaf** of the system. This sheaf was first introduced in [99] for the case of Liouville-integrable systems, but its generalization to the cases of non-Hamiltonian integrable systems and integrable systems in generalized Liouville sense is obvious.

If we restrict \mathcal{R} to the regular region \mathcal{B}_0 of \mathcal{B} (the set of regular invariant tori of the system), then \mathcal{B} is a locally trivial free Abelian sheaf of dimension p (one may view it as a \mathbb{Z}^p -bundle over \mathcal{B}_0), and its monodromy (which is a homomorphism from the fundamental group $\pi_1(\mathcal{B}_0)$ of \mathcal{B}_0 to $GL(p, \mathbb{Z})$) is nothing but the topological monodromy of the torus fibration of the regular part of the system. This topological monodromy, in the case of Liouville-integrable system, is known as the monodromy in the sense of Duistermaat [26], and it is a topological obstruction to the existence of global action-angle variables. In the case of Liouville-integrable systems with only nondegenerate elliptic singularities, studied by Boucetta and Molino [13], \mathcal{R} is still a locally free Abelian sheaf of dimension $p = \frac{1}{2} \dim M$.

When the system has non-elliptic singularities, the structure of \mathcal{R} can be quite complicated, even locally, and it contains a lot more information than the monodromy in the sense Duistermaat. For example, in the case of 2-degree-of-freedom Liouville-integrable systems restricted to isoenergy 3-manifolds, \mathcal{R} contains information on the “marks” of the so-called **Fomenko-Zieschang invariant**, which is a complete topological invariant for such systems, see e.g. [35, 11]. In fact, as found out by Fomenko, these isoenergy 3-manifolds are graph-manifolds, so the classical theory of graph-manifolds can be applied to the topological study of these 2-degree-of-freedom Liouville-integrable systems. A simple explanation of the fact that these manifolds are graph-manifolds is that they admit local \mathbb{T}^1 -actions.

The second cohomology group $H^2(\mathcal{B}, \mathcal{R})$ plays an important role in the global topological study of integrable systems, at least in the Liouville-integrable case. In fact, if two Liouville-integrable Hamiltonian systems have the same base space, the same singularities, and the same toric monodromy sheaf, then their remaining topological difference can be characterized by an element in $H^2(\mathcal{B}, \mathcal{R})$, called the (relative) **Chern class**. We refer to [99] for a precise definition of this Chern class for Liouville-integrable Hamiltonian systems (the definition is quite technical when the system has non-elliptic singularities), and the corresponding topological classification theorem. In the case of systems without singularities or with only elliptic singularities, this Chern class was first defined and studied by Duistermaat [26], and then by Dazord–Delzant [24] and Boucetta–Molino [13].

5.2. Affine base space, integrable surgery, and convexity.

In the case of Liouville-integrable systems, the base space \mathcal{B} has a natural stratified integral affine structure (local action functions of the system project to integral affine functions on \mathcal{B}), and the structure of the toric monodromy sheaf \mathcal{R} can be read off the affine structure of \mathcal{B} , see [99].

The integral affine structure on \mathcal{B} plays an important role in the problem of quantization of Liouville-integrable systems. A general idea, supported by recent works on quantization of integrable systems, see e.g. [74, 19, 69], is that one can think of Bohr-Sommerfeld or quasi-classical quantization as a discretization of the integral affine structure of \mathcal{B} : after quantization, in place of a stratified integral affine manifold, we get a “stratified nonlinear lattice” (of joint spectrum of the system). The monodromy of this joint spectrum stratified lattice of the quantized system (called *quantum monodromy*) naturally resembles the monodromy of the classical system.

The idea of **integrable surgery**, introduced in [99], is as follows: if we look at integrable systems from differential topology point of view (singular torus fibrations), instead of dynamical point of view (quasi-periodic flows), then we can perform surgery on them in order to study their properties and obtain new integrable systems from old ones. The surgery is first performed at the base space level, and then lifted to the phase space. As a side product, we also obtain new symplectic manifolds from old ones.

Let us indicate here a few interesting results obtained in [99, 95] with the help of integrable surgery:

- A very simple example of an exotic symplectic space \mathbb{R}^{2n} (which is diffeomorphic to a standard symplectic space $(\mathbb{R}^{2n}, \sum dx_i \wedge dy_i)$ but cannot be symplectically embedded into $(\mathbb{R}^{2n}, \sum dx_i \wedge dy_i)$).
- Construction of integrable systems on symplectic manifolds diffeomorphic to K3 surfaces, and on other symplectic 4-manifolds.
- Existence of a fake base space, i.e. a stratified integral affine manifold which can be realized locally as a base space of a Liouville-integrable system but globally cannot.
- A new simple proof [95] of the monodromy formula around focus-focus singularities (and their degenerate analogs) using Duistermaat–Heckman formula. The idea itself goes beyond focus-focus singularities and can be applied to other situations as well, for example in order to obtain fractional monodromy [69] via Duistermaat-Heckman formula.

Integrable surgery was recently adopted by Symington in her work on symplectic 4-manifolds [79, 80], and in the work of Leung and Symington [56] where they gave a complete list of diffeomorphism types of 4-dimensional closed “almost toric” symplectic manifolds. It seems that integrable surgery was also used by Kontsevich and Soibelman in a recent work on mirror symmetry [52].

For a noncommutatively integrable Hamiltonian system on symplectic manifolds, local action functions defined by the Mineur-Arnold formula still project to local functions on the base space \mathcal{B} , but since the number of independent action functions is equal to the dimension of invariant tori and is smaller than the dimension of \mathcal{B} , they don’t define an affine structure on \mathcal{B} , but rather an integral affine structure transverse to a singular foliation in \mathcal{B} , and under some properness condition this transverse affine structure projects to an affine structure on a quotient space $\widehat{\mathcal{B}}$ of \mathcal{B} , which has the same dimension as that of invariant tori and which may be called the **reduced affine base space**.

A particular situation where $\widehat{\mathcal{B}}$ looks nice is the case of noncommutatively integrable systems generated by Hamiltonian actions of compact Lie groups, or more generally of *proper symplectic groupoids*, see [100]. It was shown in [100] that in this case $\widehat{\mathcal{B}}$ is an integral affine manifold with locally convex polyhedral boundary, and we have a kind of **intrinsic convexity** from which one can recover various convexity theorems for momentum maps in symplectic geometry, including, for example:

- Atiyah–Guillemin–Sternberg–Kirwan convexity theorem [3, 39, 51] which says that if G is a connected compact Lie group which acts Hamiltonianly on a connected compact symplectic manifold M with an equivariant momentum map $\mu : M \rightarrow \mathfrak{g}^*$ then $\mu(M) \cap \mathfrak{t}_+^*$ is a convex polytope, where \mathfrak{t}_+^* denotes a Weyl chamber in the dual of a Cartan subalgebra of the Lie algebra of G .
- Flaschka–Ratiu’s convexity theorem for momentum maps of Poisson actions of compact Poisson-Lie groups [33].
- If one works in an even more general setting of Hamiltonian spaces of proper quasi-symplectic groupoids [89], then one also recovers Alekseev–Malkin–Meinrenken’s convexity theorem for group-valued momentum maps [2].

In order to further incite the reader to read [100], let us mention here a recent beautiful convexity theorem of Weinstein [86] which has a clear meaning in Hamiltonian dynamics and which fits well in the above framework of proper symplectic groupoid actions and noncommutatively integrable Hamiltonian systems:

Theorem 5.1 (Weinstein [86]). *For any positive-definite quadratic Hamiltonian function H on the standard symplectic space \mathbb{R}^{2k} , denote by $\phi(H)$ the k -tuple $\lambda_1 \leq \dots \leq \lambda_k$ of frequencies of H ordered non-decreasingly, i.e. H can be written as $H = \sum \lambda_i(x_i^2 + y_i^2)$ in a canonical coordinate system. Then for any two given positive nondecreasing n -tuples $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\gamma = (\gamma_1, \dots, \gamma_k)$, the set*

$$(5.1) \quad \Phi_{\lambda, \gamma} = \{\phi(H_1 + H_2) \mid \phi(H_1) = \lambda, \phi(H_2) = \gamma\}$$

is a closed, convex, locally polyhedral subset of \mathbb{R}^k .

Note the above set $\Phi_{\lambda, \gamma}$ is closed but not bounded. For example, when $k = 1$ then $\Phi_{\lambda, \gamma}$ is a half-line.

5.3. Localization formulas.

A general idea in analysis and geometry is to express global invariants in terms of local invariants, via *localization formulas*.

Various global topological invariants, including the Chern classes (of the tangent bundle), of the symplectic ambient manifold of a Liouville-integrable system, can be localized at singularities of the system. Some results in this direction can be found in recent papers of Gross [38] and Smith [77], though much still waits to be worked out for general integrable systems. For example, consider a 4-dimensional symplectic manifold with a proper integrable system whose fixed points are nondegenerate. Then to find c_2 (the Euler class) of the manifold, one simply needs to count the number of fixed points with signs: the plus sign for elliptic-elliptic ($k_e = 2, k_h = k_f = 0$ in Williamson type), hyperbolic-hyperbolic ($k_h = 2$) and focus-focus points, and the minus sign for elliptic-hyperbolic ($k_e = k_h = 1$) points.

In symplectic geometry, there is a famous localization formula for Hamiltonian torus actions, due to Duistermaat and Heckman [28]. There is a topological version of this formula, in terms of equivariant cohomology, due to Atiyah–Bott [4] and Berline–Vergne [8], and a non-Abelian version due to Witten [88] and Jeffrey–Kirwan [45]. We refer to [5, 27, 41] for an introduction to these formulas. It would be nice to have analogs of these formulas for proper integrable systems.

6. INFINITE-DIMENSIONAL TORUS ACTIONS

A general idea is that proper infinite dimensional integrable systems admit infinite-dimensional torus actions. Consider, for example, the KdV equation

$$(6.1) \quad u_t = -u_{xxx} + 6uu_x$$

with periodic boundary condition $u(t, x + 1) = u(t, x)$. We will view this KdV equation as a flow on the space of functions u on $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$. Then it is a Hamiltonian equation with the Hamiltonian

$$(6.2) \quad H = \int_{\mathbb{S}^1} \left(\frac{1}{2} u_x^2 + u^3 \right) dx$$

and the Poisson structure $\frac{d}{dx}$, see e.g. [49]. In other words, if F and G are two functional on the space of functions on \mathbb{S}^1 then their Poisson bracket is

$$(6.3) \quad \{F, G\} = \int_{\mathbb{S}^1} \frac{\partial F}{\partial u} \frac{d}{dx} \frac{\partial G}{\partial u} dx .$$

The Poisson structure $\frac{d}{dx}$ admits a Casimir function

$$(6.4) \quad [u] = \int_{\mathbb{S}^1} u dx .$$

The Sobolev space

$$(6.5) \quad \mathcal{H}_0^1 = \left\{ u : \mathbb{S}^1 \rightarrow \mathbb{R}^1 \mid \int_{\mathbb{S}^1} (u_x^2 + u^2) dx < \infty, [u] = 0 \right\}$$

together with the Poisson structure $\frac{d}{dx}$ is a *weak* symplectic infinite-dimensional manifold, which is symplectomorphic, via the Fourier transform, to the symplectic Hilbert space

$$(6.6) \quad \mathfrak{h}_{3/2} := \left\{ (x_n, y_n)_{n \in \mathbb{N}} \mid x_n, y_n \in \mathbb{R}, \sum n^3 x_n^2 + \sum n^3 y_n^2 < \infty \right\}$$

with the *weak* symplectic structure $\omega = \sum dx_n \wedge dy_n$. We have the following Birkhoff normal form theorem for the periodic KdV equation, due to Kappeler and his collaborators Bättig, Bloch, Guillot, Mityagin, Makarov, see [16, 49] and references therein:

Theorem 6.1 (Kappeler et al.). *There is a bi-analytic 1-1 symplectomorphism $\Psi : \mathfrak{h}_{3/2} \rightarrow \mathcal{H}_0^1$, such that the coordinates (x, y) become global Birkhoff coordinates for the KdV equation under this symplectomorphism, i.e. the transformed Hamiltonian $H_\Psi = H \circ \Psi$ depend only on $x_n^2 + y_n^2, n \in \mathbb{N}$.*

See [49] for a more precise and general statement of the above theorem. A direct consequence of Theorem 6.1 is that we have a Hamiltonian infinite-dimensional torus action generated by the action functions $I_n = x_n^2 + y_n^2$ which preserve the KdV system and whose orbits are exactly the level sets of the KdV. In fact, these action functions are also found by the Mineur-Arnold integral formula.

Remark that the level sets $N_c := \{I_n = c_n \forall n \in \mathbb{N}\}$, where c_n are nonnegative constants such that $\sum n^3 c_n < \infty$, are *compact* with respect to the induced norm topology. They are *not* submanifolds of the phase space, but can still be viewed as (infinite-dimensional) Liouville tori. The natural topology that we have to put on the infinite-dimensional torus is the *product topology*, and then it becomes a compact topological group by Tikhonoff theorem, and the Hamiltonian action $\mathbb{T}^\infty \times \mathcal{H}_0^1 \rightarrow \mathcal{H}_0^1$ is a continuous action.

So basically the periodic KdV is just an infinite-dimensional oscillator. The periodic defocusing NLS (non-linear Schrödinger) equation is similar and is also an infinite-dimensional oscillator, see [16, 37].

There are other integrable equations, like the sine-Gordon equation and the focusing NLS equation, which are topologically very different from infinite-dimensional oscillators: they admit *unstable* singularities. Their topological structure was studied to some extent by McKean, Ercolani, Forest, McLaughlin and many other people, see e.g. [62, 31, 57] and references therein.

A general idea is that, even when an integrable PDE admits unstable phenomena, locally near each level set the system can be decomposed into 2 parts: an unstable part which is finite-dimensional, and an infinite-dimensional oscillator. The reason is that the energy is finite, and to get something unstable one needs big energy, so one can only get finitely many unstable things, the rest is just a small (infinite-dimensional) oscillator. This idea reduces the topological study of infinite-dimensional integrable systems to that of finite-dimensional systems.

A concrete case study, namely the symplectic topology of the focusing NLS equation,

$$(6.7) \quad -iq_t = q_{xx} + 2\bar{q}q^2 ,$$

where q is a complex-valued function on \mathbb{S}^1 for each t , is the subject of a joint work in progress of Thomas Kappeler, Peter Topalov and myself [50]. It is a Hamiltonian system on the Sobolev space $\mathcal{H}^1(\mathbb{S}^1, \mathbb{C})$ of complex-valued functions on the circle with the Poisson bracket

$$(6.8) \quad \{F, G\} = i \int_{\mathbb{S}^1} \left(\frac{\partial F}{\partial q} \frac{\partial G}{\partial \bar{q}} - \frac{\partial G}{\partial q} \frac{\partial F}{\partial \bar{q}} \right) dx ,$$

and its Hamiltonian is

$$(6.9) \quad H = \int_{\mathbb{S}^1} (q_x \bar{q}_x - q^2 \bar{q}^2) dx .$$

Here is our speculation as to what happens there: For any q which belongs to a dense open subset \mathcal{M}_0 of $\mathcal{H}^1(\mathbb{S}^1, \mathbb{C})$ (the “almost-regular set”), the level set $Iso(q)$ (which is the same as the isospectral set of the Zakharov-Shabat operator) is a torus (of infinite dimension in general), and there is a neighborhood $\mathcal{U}(Iso(q))$ of $Iso(q)$ in the phase space which admits a full action-angle system of coordinates similar to the KdV case. For any $q \notin \mathcal{M}_0$, a neighborhood $\mathcal{U}(Iso(q))$ of $Iso(q)$ still admits a torus action of “finite corank” and we have a partial Birkhoff coordinate system of finite codimension. If $q \notin \mathcal{M}_0$ is “nondegenerate” (it corresponds to a condition on the spectrum of q), then $\mathcal{U}(Iso(q))$ together with the fibration by the level sets can be written topologically as a direct product of an infinite-dimensional oscillator and a finite number of focus-focus singularities. The fact that unstable nondegenerate singularities of the focusing NLS are of focus-focus type can be seen from the work of Li and McLaughlin [57], and is probably due to the \mathbb{T}^1 -symmetry of the system (translations in x -variable). If we restrict the focusing NLS system to even functions ($q(-x) = q(x)$), then it is still an integrable Hamiltonian system, but with hyperbolic instead of focus-focus singularities.

REFERENCES

- [1] M. Adler, P. van Moerbeke, and P. Vanhaecke, *Algebraic completely integrable systems, Painlevé architecture and Lie algebras*, Springer, 2004.
- [2] Anton Alekseev, Anton Malkin, and Eckhard Meinrenken, *Lie group valued moment maps*, J. Differ. Geom. **48** (1998), no. 3, 445–495.
- [3] M. F. Atiyah, *Convexity and commuting Hamiltonians*, Bull. London Math. Soc. **14** (1982), no. 1, 1–15.
- [4] M. F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology **23** (1984), no. 1, 1–28.
- [5] Michèle Audin, *The topology of torus actions on symplectic manifolds*, Birkhäuser Verlag, Basel, 1991.

- [6] Michèle Audin, *Hamiltonian monodromy via Picard-Lefschetz theory*, Commun. Math. Phys. **229** (2002), no. 3, 459–489.
- [7] Larry Bates and Richard Cushman, *What is a completely integrable nonholonomic dynamical system ?*, Reports Math. Phys. **44** (1999), no. 1-2, 29–35.
- [8] N. Berline and M. Vergne, *Classes caractéristiques équivariantes. formule de localisation en cohomologie équivariante*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 9, 539–541.
- [9] Salomon Bochner, *Compact groups of differentiable transformations*, Annals of Math. (2) **46** (1945), 372–381.
- [10] O.I. Bogoyavlenskij, *Extended integrability and bi-hamiltonian systems*, Comm. Math. Phys. **196** (1998), no. 1, 19–51.
- [11] A.V. Bolsinov and A.T. Fomenko, *Integrable Hamiltonian Systems. Geometry, Topology, Classification. Vol. 1 and 2 (in Russian)*, 1999.
- [12] A.V. Bolsinov and B. Jovanovic, *Non-commutative integrability, moment map and geodesic flows*, Ann. Global Anal. Geom. **23** (2003), no. 4, 305–322.
- [13] M. Boucetta and P. Molino, *Géométrie globale des systèmes hamiltoniens complètement intégrables*, C. R. Acad. Sci. Paris, Ser. I **308** (1989), 421–424.
- [14] A. D. Bruno, *Local methods in nonlinear differential equations*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1989.
- [15] A.D. Bruno and S. Walcher, *Symmetries and convergence of normalizing transformations*, J. Math. Anal. Appl. **183** (1994), 571–576.
- [16] D. Bättig, A.M. Bloch, J.C. Guillot, and T. Kappeler, *On the symplectic structure of the phase space for periodic KdV, Toda, and defocusing NLS*, Duke Math. J. **79** (1995), no. 3, 549–604.
- [17] Yves Colin de Verdière, *Singular Lagrangian manifolds and semiclassical analysis*, Duke Math. J. **116** (2003), no. 2, 263–298.
- [18] Yves Colin de Verdière and J. Vey, *Le lemme de Morse isochore*, Topology **18** (1979), 283–293.
- [19] Yves Colin de Verdière and San Vu-Ngoc, *Singular Bohr-Sommerfeld rules for 2D integrable systems*, Ann. Sci. Ecole Norm. Sup. (4) **36** (2003), 1–55.
- [20] M. Condevaux, P. Dazord, and P. Molino, *Géométrie du moment*, Séminaire Sud-Rhodanien I, Publications du département de math., Univ. Claude Bernard - Lyon I (1988), 131–160.
- [21] Carlos Currás Bosch and Eva Miranda, *Symplectic linearization of singular Lagrangian foliations in M^4* , Diff. Geom. Appl. **18** (2003), no. 2, 195–205.
- [22] Richard Cushman and Hans Duistermaat, *Non-Hamiltonian monodromy*, J. Diff. Equations **172** (2001), no. 1, 42–58.
- [23] Richard Cushman and Reyer Sjamaar, *On singular reduction of Hamiltonian spaces*, Symplectic geometry and mathematical physics (Aix-en-Provence, 1990), Progr. Math., 99, Birkhäuser, 1991, pp. 114–128.
- [24] Pierre Dazord and Thomas Delzant, *Le problème général des variables action-angles*, J. Diff. Geom. **26** (1987), no. 2, 223–251.
- [25] J.-P. Dufour and P. Molino, *Compactification d’actions de \mathbb{R}^n et variables action-angle avec singularités*, Symplectic geometry, groupoids, and integrable systems (Berkeley, CA, 1989), Springer, New York, 1991, pp. 151–167.
- [26] Hans Duistermaat, *On global action-angle variables*, Comm. Pure Appl. Math. **33** (1980), 687–706.
- [27] ———, *Equivariant cohomology and stationary phase*, Symplectic geometry and quantization (Sanda and Yokohama, 1993), Contemp. Math., 179, Amer. Math. Soc., 1994, pp. 45–62.
- [28] Hans Duistermaat and G.H. Heckman, *On the variation in the cohomology of the symplectic form of the reduced phase space*, Invent. Math. **69** (1982), 259–269.
- [29] L. H. Eliasson, *Normal forms for Hamiltonian systems with Poisson commuting integrals*, Ph.D. Thesis (1984).
- [30] ———, *Normal forms for Hamiltonian systems with Poisson commuting integrals—elliptic case*, Comment. Math. Helv. **65** (1990), no. 1, 4–35.
- [31] N.M. Ercolani and D.W. McLaughlin, *Toward a topological classification of integrable PDE’s*, The Geometry of Hamiltonian Systems. MSRI Publ. Vol. 22 (1991), 111–130.
- [32] N.W. Evans, *Superintegrability of the Calogero–Moser system*, Phys. Lett. A **95** (1983), 279.
- [33] Hermann Flaschka and Tudor Ratiu, *A convexity theorem for Poisson actions of compact Lie groups*, Ann. Sci. École Norm. Sup. (4) **29** (1996), no. 6, 787–809.

- [34] A.T. Fomenko, *Integrability and nonintegrability in geometry and mechanics*, Kluwer, Dordrecht, 1988.
- [35] A.T. Fomenko and H. Zieschang, *A topological invariant and a criterion for the equivalence of integrable hamiltonian systems with two degrees of freedom*, Math. USSR-Izv. **36** (1991), no. 3, 567–596.
- [36] Jean-Pierre Francoise, *Intégrales de périodes en géométries symplectique et isochore*, Géométrie symplectique et mécanique, Colloq. Int. Sémin. Sud- Rhodan. Géom. V, La Grande Motte 1988, Lect. Notes Math. 1416, 105-138 , 1990.
- [37] B Grébert, T. Kapeler, and J. Pöschel, *Normal form theory for the NLS equation*, preprint (2003).
- [38] M. Gross, *Topological mirror symmetry*, Invent. Math. **144** (2001), 75–137.
- [39] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), 491–513.
- [40] ———, *On collective complete integrability according to the method of Thimm*, Ergodic Theory Dynam. Systems **3** (1983), no. 2, 219–230.
- [41] Victor Guillemin, Viktor Ginzburg, and Yael Karshon, *Moment maps, cobordisms, and Hamiltonian group actions*, Mathematical Surveys and Monographs, vol. 98, American Mathematical Society, Providence, RI, 2002.
- [42] Hidekazu Ito, *Convergence of Birkhoff normal forms for integrable systems*, Comment. Math. Helv. **64** (1989), no. 3, 412–461.
- [43] ———, *Action-angle coordinates at singularities for analytic integrable systems*, Math. Z. **206** (1991), no. 3, 363–407.
- [44] ———, *Integrability of Hamiltonian systems and Birkhoff normal forms in the simple resonance case*, Math. Ann. **292** (1992), no. 3, 411–444.
- [45] L. Jeffrey and F. Kirwan, *Localization for nonabelian group actions*, Topology **34** (1995), no. 2, 291–327.
- [46] Bozidar Jovanovic, *On the integrability of geodesic flows of submersion metrics*, Lett. Math. Phys. **61** (2002), no. 1, 29–39.
- [47] V. Kalashnikov, *Generic integrable Hamiltonian systems on a four-dimensional symplectic manifold*, Izv. Math. **62** (1998), no. 2, 261–285.
- [48] T. Kappeler, Y. Kodama, and A. Némethi, *On the Birkhoff normal form of a completely integrable Hamiltonian system near a fixed point with resonance*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **XXVI** (1998), no. 4, 623–661.
- [49] T. Kappeler and J. Pöschel, *KdV and KAM*, Springer, 2003.
- [50] T. Kappeler, T. Topalov, and N.T. Zung, *Symplectic topology of the focusing NLS equation*, work in progress (2004).
- [51] Frances Kirwan, *Convexity properties of the moment mapping. III*, Invent. Math. **77** (1984), no. 3, 547–552.
- [52] M. Kontsevich and Y. Soibelman, *Affine structure and non-archimedean analytic spaces*, preprint math.AG/0406564 (2004).
- [53] Jeroen S. W. Lamb and John A. G. Roberts, *Time-reversal symmetry in dynamical systems: a survey*, Phys. D **112** (1998), no. 1-2, 1–39.
- [54] L.M. Lerman and Ya.L. Umanskij, *Structure of the Poisson action of \mathbb{R}^2 on a four-dimensional symplectic manifold. I*, Selecta Math. Sov. **6** (1987), 365–396.
- [55] ———, *Structure of the Poisson action of \mathbb{R}^2 on a four-dimensional symplectic manifold. II*, Selecta Math. Sov. **7** (1988), 39–48.
- [56] N.C. Leung and M. Symington, *Almost toric symplectic four-manifolds*, preprint (2003).
- [57] Y. Li and D. McLaughlin, *Morse and Melnikov functions for NLS Pde's*, Comm. Math. Phys. **162** (1994), 175–214.
- [58] Joseph Liouville, *Note sur l'intégration des équations différentielles de la dynamique, présentée au bureau des longitudes le 29 juin 1853*, Journal de Math. Pures et Appl. **20** (1855), 137–138.
- [59] S. Lojasiewicz, *Sur le problème de la division*, Studia Math. **18** (1959), 87–136.
- [60] M. M. Golubitsky and I. I. Stewart, *Generic bifurcation of Hamiltonian systems with symmetry*, Physica D **24** (1987), 391–405.
- [61] Y. Matsumoto, *Topology of torus fibrations*, Sugaku Expositions **2** (1989), 55–73.
- [62] H.P. McKean, *The sine-Gordon and sinh-Gordon equations on the circle*, Comm. Pure Appl. Math. **34** (1981), 197–257.

- [63] Henri Mineur, *Sur les systemes mecaniques admettant n integrales premieres uniformes et l'extension a ces systemes de la methode de quantification de Sommerfeld*, C. R. Acad. Sci., Paris **200** (1935), 1571–1573 (French).
- [64] ———, *Sur les systemes mecaniques dans lesquels figurent des parametres fonctions du temps. Etude des systemes admettant n integrales premieres uniformes en involution. Extension a ces systemes des conditions de quantification de Bohr-Sommerfeld.*, Journal de l'Ecole Polytechnique, Série III, 143ème année (1937), 173–191 and 237–270.
- [65] Eva Miranda and Vu Ngoc San, *A singular Poincaré lemma*, preprint math.SG/0405430 (2004).
- [66] Eva Miranda and Nguyen Tien Zung, *Equivariant normal form for nondegenerate singular orbits of integrable Hamiltonian systems*, preprint 2003, submitted to Annales Ecole Norm. Sup. (2004).
- [67] A. S. Mischenko and A. T. Fomenko, *A generalized method for Liouville integration of Hamiltonian systems*, Funct. Anal. Appl. **12** (1978), 46–56.
- [68] N.N. Nekhoroshev, *Action-angle variables and their generalizations*, Trans. Moscow Math. Soc. **26** (1972), 180–198.
- [69] N.N. Nekhoroshev, D. Sadovskii, and B. Zhilinskii, *Fractional monodromy of resonant classical and quantum oscillators*, C. R. Acad. Sci. Paris Ser. I **335** (2002), no. 11, 985–988.
- [70] Valentin Poënar, *Singularités C^∞ en présence de symétrie*, Lecture Notes in Mathematics, Vol. 510, 1976.
- [71] N. Reshetikhin, *Integrability of characteristic Hamiltonian systems on simple Lie groups with standard Poisson Lie structure*, Comm. Math. Phys. **242** (2003), 1–29.
- [72] Robert Roussarie, *Modèles locaux de champs et de formes*, Astérisque, Société Mathématique de France, Paris, 1975.
- [73] H. Rüssmann, *Über das Verhalten analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtslösung*, Math. Ann. **154** (1964), 285–300.
- [74] Vu Ngoc San, *Bohr-Sommerfeld conditions for integrable systems with critical manifolds of focus-focus type*, Comm. Pure Appl. Math. **53** (2000), no. 2, 143–217.
- [75] ———, *Formes normales semi-classiques des systèmes complètement intégrables au voisinage d'un point critique de l'application moment*, Asymptotic Anal. **24** (2000), no. 3–4, 319–342.
- [76] ———, *Symplectic techniques for semiclassical completely integrable systems*, This Volume (2004).
- [77] Ivan Smith, *Torus fibrations on symplectic four-manifolds*, Turkish J. Math. **25** (2001), no. 1, 69–95.
- [78] Laurent Stolovitch, *Singular complete integrability*, Publications IHES **91** (2003), 134–210.
- [79] Margaret Symington, *Generalized symplectic rational blowdown*, Algebr. Geom. Topol. **1** (2001), 503–518.
- [80] ———, *Four dimensions from two in symplectic topology*, preprint math.SG/0210033 (2002).
- [81] A.P. Veselov and S.P. Novikov, *Poisson brackets and complex tori*, Proceedings of Steklov Institute of Mathematics **165** (1984), no. 3, 53–65.
- [82] Jacques Vey, *Sur certains systèmes dynamiques séparables*, Amer. J. Math. **100** (1978), no. 3, 591–614.
- [83] ———, *Algèbres commutatives de champs de vecteurs isochores*, Bull. Soc. Math. France **107** (1979), 423–432.
- [84] G. Wassermann, *Classification of singularities with compact Abelian symmetry*, Banach Center Publications **20** (1988), 475–498.
- [85] Alan Weinstein, *Poisson geometry*, Differential Geom. Appl. **9** (1998), no. 1-2, 213–238.
- [86] ———, *Poisson geometry of discrete series orbits, and momentum convexity for noncompact group actions.*, Lett. Math. Phys. **56** (2001), no. 1, 17–30.
- [87] J. Williamson, *On the algebraic problem concerning the normal forms of linear dynamical systems*, Amer. J. Math. **58** (1936), no. 1, 141–163.
- [88] E. Witten, *Two-dimensional gauge theories revisited*, J. Geom. Phys. **9** (1992), no. 4, 303–368.
- [89] Ping Xu, *Momentum maps and Morita equivalence*, preprint math.SG/0307319 (2003).
- [90] M. Zhitomirskii, *Normal forms of symmetric Hamiltonian systems*, J. Differential Equations **111** (1994), no. 1, 58–78.

- [91] Nguyen Tien Zung, *On the general position property of simple Bott integrals*, Russ. Math. Surv. **45** (1990), no. 4, 179–180.
- [92] ———, *Symplectic topology of integrable Hamiltonian systems. I: Arnold-Liouville with singularities*, Compos. Math. **101** (1996), no. 2, 179–215.
- [93] ———, *A note on degenerate corank-one singularities of integrable Hamiltonian systems*, Comment. Math. Helv. **75** (2000), no. 2, 271–283.
- [94] ———, *Convergence versus integrability in Birkhoff normal form*, preprint math.DS/0104279, to appear in Annals Math. (2001).
- [95] ———, *Another note on focus-focus singularities*, Lett. Math. Phys. **60** (2002), no. 1, 87–99.
- [96] ———, *Convergence versus integrability in Poincaré-Dulac normal form*, Math. Res. Lett. **9** (2002), no. 2-3, 217–228.
- [97] ———, *Reduction and integrability*, preprint arxiv math.DS/0201087 (2002).
- [98] ———, *Actions toriques et groupes d'automorphismes de singularités de systèmes dynamiques intégrables*, C. R. Acad. Sci. Paris **336** (2003), no. 12, 1015–1020.
- [99] ———, *Symplectic topology of integrable Hamiltonian systems. II: Topological classification*, Compositio Math. **138** (2003), no. 2, 125–156.
- [100] ———, *Proper groupoids and momentum maps: linearization, affinity, and convexity*, preprint math.SG/0407208 (2004).

LABORATOIRE EMILE PICARD, UMR 5580 CNRS, UFR MIG, UNIVERSITÉ TOULOUSE III

E-mail address: `tienzung@picard.ups-tlse.fr`